

Chronic Adaptations to Eccentric Training: A Systematic Review

Jamie Douglas^{1,2} · Simon Pearson^{1,3} · Angus Ross² · Mike McGuigan^{1,4}

© Springer International Publishing Switzerland 2016

Abstract

Background Resistance training is an integral component of physical preparation for athletes. A growing body of evidence indicates that eccentric strength training methods induce novel stimuli for neuromuscular adaptations.

Objective The purpose of this systematic review was to determine the effects of eccentric training in comparison to concentric-only or traditional (i.e. constrained by concentric strength) resistance training.

Methods Searches were performed using the electronic databases MEDLINE via EBSCO, PubMed and SPORTDiscus via EBSCO. Full journal articles investigating the long-term (≥ 4 weeks) effects of eccentric training in healthy (absence of injury or illness during the 4 weeks preceding the training intervention), adult (17–35 years), human participants were selected for the systematic review. A total of 40 studies conformed to these criteria.

Results Eccentric training elicits greater improvements in muscle strength, although in a largely mode-specific manner. Superior enhancements in power and stretch-shortening cycle (SSC) function have also been reported. Eccentric training is at least as effective as other modalities

in increasing muscle cross-sectional area (CSA), while the pattern of hypertrophy appears nuanced and increased CSA may occur longitudinally within muscle (i.e. the addition of sarcomeres in series). There appears to be a preferential increase in the size of type II muscle fibres and the potential to exert a unique effect upon fibre type transitions. Qualitative and quantitative changes in tendon tissue that may be related to the magnitude of strain imposed have also been reported with eccentric training.

Conclusions Eccentric training is a potent stimulus for enhancements in muscle mechanical function, and muscle-tendon unit (MTU) morphological and architectural adaptations. The inclusion of eccentric loads not constrained by concentric strength appears to be superior to traditional resistance training in improving variables associated with strength, power and speed performance.

Key Points

Eccentric training can improve muscle mechanical function to a greater extent than other modalities.

Novel muscle-tendon unit adaptations associated with a faster (i.e. explosive) phenotype have been reported.

Eccentric training may be especially beneficial in enhancing strength, power and speed performance.

✉ Jamie Douglas
jamie.douglas@hpsnz.org.nz

¹ Sports Performance Research Institute New Zealand (SPRINZ), Auckland University of Technology, Auckland, New Zealand

² High Performance Sport New Zealand (HPSNZ), AUT Millennium, 17 Antares Place, Mairangi Bay, Auckland 0632, New Zealand

³ Queensland Academy of Sport, Nathan, QLD, Australia

⁴ School of Medical and Health Sciences, Edith Cowan University, Perth, WA, Australia

1 Background

Resistance training has become a ubiquitous component of physical preparation programmes for athletic populations [1]. It has been well established that resistance training can

improve a host of neuromuscular variables relevant to athletic performance across a continuum of strength, power, and endurance events [1–3]. Traditional resistance training typically includes both eccentric and concentric phases of movement across a set of repetitions. Eccentric muscle actions occur when the load applied to the muscle exceeds the force produced by the muscle itself, resulting in a lengthening action [4]. Therefore, muscle forces tend to be highest during lengthening actions [5]. The prescription of load is dictated by concentric strength and thus tends to insufficiently load the eccentric phase of movement. A growing body of evidence indicates that resistance training programmes that sufficiently load the eccentric phase of movement can elicit superior neuromuscular adaptations compared with concentric-only or traditional resistance training constrained by concentric strength [6–8]. The training stress and physiological strain imposed by eccentric training induces an adaptive response conducive to enhancements in muscle mechanical function, and alterations in muscle-tendon unit (MTU) morphology and architecture. Metrics of strength, power and stretch-shortening cycle (SSC) function appear to be particularly responsive to eccentric stimuli. The purpose of this review was to systematically retrieve and collate studies that directly compared eccentric training with concentric or traditional resistance training.

2 Methods

The review was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines for systematic reviews [9]. One reviewer performed initial database searches for articles investigating the chronic (i.e. ≥ 4 weeks) adaptations to eccentric training interventions on in vivo muscle-tendon properties and performance in human subjects (last search April 2016). Searches were performed using the electronic databases MEDLINE via EBSCO (1950–present), PubMed (1950–present) and SPORTDiscus via EBSCO (1985–present). Key search terms were grouped and searched within the article title, abstract, and keywords using the search conjunctions ‘OR’ and ‘AND’. Combinations of the following terms were used as search terms: ‘eccentric exercise’, ‘eccentric training’, ‘eccentric contraction’, ‘lengthening contraction’, ‘negative work’ and ‘passive work’ in conjunction with the terms ‘muscle’, ‘tendon’, ‘strength’, ‘power’, ‘speed’, ‘hypertrophy’, ‘force’, ‘velocity’ and ‘performance’. Key journals identified were also searched using the keyword ‘eccentric’. Furthermore, the reference lists of articles retrieved were screened for additional eligible articles. Full journal articles investigating the long-term effects of eccentric training (≥ 4 weeks)

in healthy (i.e. the absence of injury or illness during the 4 weeks preceding the training intervention), adult (i.e. 17–35 years), human participants were selected for systematic review (Fig. 1). Articles were excluded if the aforementioned criteria were not fulfilled, training was performed less than twice weekly, eccentric exercise intensity was not quantified or was below the relative concentric exercise intensity, or no concentric or traditional resistance training control group was included.


3 Results and Discussion

3.1 Participant and Intervention Characteristics

Across the 40 studies included for review, 1150 participants (406 females and 744 males) with a mean age of 23.9 years (range 17.6–35.0) were recruited. The majority of investigations (32/40; 80 %) recruited untrained participants, four (10 %) recruited participants with resistance training experience (3 months to 1 year), and the remaining four (10 %) recruited participants who were either moderately trained or participated in elite sport. The majority of investigations compared eccentric training (i.e. eccentric contractions at an intensity above the relative concentric training intensity, performed alone or in conjunction with concentric contractions) of various volumes and intensities with traditional resistance training (i.e. mixed eccentric and concentric contractions limited by concentric intensity) and/or concentric training (i.e. concentric contractions only). A non-training control group was included in 17 (43 %) studies. Other training variables compared included the magnitude of overload (i.e. intensity; heavy vs. light), contraction velocity (i.e. tempo; fast vs. slow) and the additional effects of whey protein hydrolysate supplementation. The average intervention duration was 9.8 weeks (range 5–20), with a training frequency of 2.9 sessions per week (range 2.0–4.2). Single-joint movements were predominantly investigated (32 studies; 80 %) and isokinetic modalities were used more (26 studies; 65 %) than isoinertial modalities.

3.2 Muscle Mechanical Function

Eccentric training has been consistently reported to increase concentric [10–32], isometric [18, 23, 25, 29, 33–37], and eccentric [11, 13–15, 17, 18, 21–25, 27–29, 31, 37–42] strength when assessed via isoinertial (i.e. repetition maximum [RM] testing) or isokinetic (i.e. maximal voluntary contraction [MVC] testing) modalities (Table 1). Concentric training also elicits increases in concentric [11, 13–18, 21–32, 37–41], isometric [18, 23, 29, 33, 37, 40] and eccentric

Fig. 1 Search strategy

[11, 13–15, 17, 18, 21, 23, 25, 27–29, 39, 42] strength, while increases in concentric [10, 12, 19, 20, 23], isometric [23, 34] and eccentric strength [23] are similarly observed following traditional resistance training (Table 1). Strength increases have been proposed to be largely mode-specific [8], and while some studies reported that eccentric training increased eccentric strength to a greater extent compared with concentric training [13, 21, 22, 25, 27, 31, 38, 39, 41], and vice versa [11, 38–41], others found no differences between modalities [10, 12, 14, 15, 19, 20, 23, 24, 28–30, 32–34, 42, 43]. A number of studies investigating eccentric training included the concentric portion of the movement in addition to the overloaded eccentric portion [10, 12–14, 19, 20, 23, 24, 32, 34, 42, 43], which may partly explain the mixed findings compared with previous reviews that compared eccentric-only with concentric-only modalities [6, 8]. When using eccentric loads greater than maximal concentric strength (e.g. 1RM or MVC), eccentric training generally leads to greater overall strength increases (i.e. combined concentric, isometric and eccentric strength) than concentric and traditional training [12, 16, 17, 21–23, 25–27, 30, 31]. Furthermore, studies directly comparing heavier with lighter eccentric loads found that heavier eccentric training induced greater increases in eccentric strength [16, 26]. Muscle contraction velocity used within training can also influence strength adaptations, and greater increases in eccentric strength have been observed with fast versus slow eccentric training [17],

while increases in eccentric strength with eccentric training become more pronounced when the testing velocity corresponds to that used within training [8]. Greater increases in contralateral eccentric strength (i.e. cross-education) have been reported with fast (i.e. 180°/s) versus slow (i.e. 30°/s) eccentric training [44], although improvements can also occur following training at moderate (i.e. 60°/s) contraction speeds [45]. Fast contractions have also been proposed to allow for a greater transfer of eccentric training to concentric strength [7].

Compared with changes in muscle strength, relatively few studies investigated changes in muscle power [13, 14, 19, 34, 46, 47] or contractile rate of force development (RFD) [27, 33]. Muscle power, as assessed primarily by lower body jump variations, increased with eccentric training within a number of studies, while concentric or traditional training had no clear effect [13, 34, 46, 47]. Furthermore, the finding of Colliander and Tesch [14], where vertical jump increased following concentric, but not eccentric, training, may have been a statistically spurious observation. Closer inspection of their data indicates that eccentric training was, at least practically, superior to concentric training (i.e. 8 vs. 4%; Cohen's d 0.36). Vertical jump performance involves an SSC component, and variables associated with SSC performance appear to improve to a greater extent with eccentric training. SSC efficiency (i.e. taken as the ratio of countermovement to squat jump performance), drop jump

Table 1 Studies comparing the effects of eccentric-overload and coupled maximal eccentric and concentric contractions with traditional and concentric-only resistance training on muscle mechanical function

Study, year	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Barstow et al. [43], 2003	ECC $n = 13$, TRAD $n = 13$, CONC $n = 13$ (8F, 31M); mean age: 22.2 years; training status: 3 months' resistance training	EF (isoinertial; single-joint)	Volume: 3 sets of 8 repetitions; intensity: ECC group: 100/60 % IRM; TRAD group: 60 % IRM; tempo: 2 s ECC, 2 s CONC	12 weeks (2 sessions per week)	No differences in CONC EF strength (IRM) between ECC (16 %), TRAD (14 %) or CONC (10 %)
Ben-Sira et al. [10], 1995	ECC _{Heavy} $n = 8$, TRAD $n = 8$, ECC _{Light} $n = 10$, CONC $n = 12$, CONC $n = 10$ (48F); mean age: 21.1 years; training status: untrained	KE (isoinertial; single-joint)	Volume: ECC _{Heavy} group: 3 sets of 5 repetitions, TRAD, ECC _{Light} and CONC group: 3 sets of 10 repetitions; intensity: ECC _{Heavy} group: 135/65 % IRM; TRAD, ECC _{Light} and CONC groups: 65 % IRM; tempo: 3 s ECC, 1 s CONC	8 weeks (2 sessions per week)	Increase in CONC KE strength (IRM) with ECC _{Heavy} (23 %) and TRAD (19 %) vs. CONC (4 %), no difference between interventions
Blazevich et al. [11], 2007	ECC $n = 11$, CONC $n = 10$, CONC $n = 9$ (16F, 14 M); mean age: 22.8 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 5 sets of 6 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 30°/s	10 weeks (3 sessions per week)	Increase in CONC KE strength (MVC) with ECC (16 %) and CONC (24 %) vs. CONC (1 %), but greater increase with CONC Increase in ECC KE strength (MVC) with ECC (39 %) and CONC (36 %) vs. CONC (3 %), no difference between interventions
Blazevich et al. [33], 2008	ECC $n = 11$, CONC $n = 10$ (11F, 10 M); mean age: 22.8 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 5 sets of 6 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 30°/s	10 weeks (3 sessions per week)	Increase in ISO KE strength (MVC) with ECC (10 %) and CONC (13 %), no difference between interventions Increase in RFD 30 ms ($N \cdot s^{-1}$) with ECC (28 %) and CONC (50 %), greater increase with CONC vs. ECC
Brandenberg and Docherty [12], 2002	ECC $n = 8$, TRAD $n = 10$ (18 M); mean age: NR, university students; training status: 1 year resistance training	EF and EE (isoinertial; single-joint)	Volume: ECC group: 3 sets of 10 repetitions, TRAD group: 4 sets of 10 repetitions; intensity: ECC group: 115/75 % IRM, TRAD group: 75 % IRM; tempo: 2 s ECC, 2 s CONC	9 weeks (2.8 sessions per week)	Increase in CONC EF strength (IRM) with ECC (9 %) and TRAD (11 %), no difference between interventions Increase in CONC EE strength (IRM) with ECC (24 %) and TRAD (15 %), but greater increase with ECC training
Colliander and Tesch [13], 1990	ECC + CONC $n = 11$, CONC $n = 11$, CONC $n = 7$ (29 M); mean age: 26.3 years; training status: untrained	KE (isokinetic; single-joint)	Volume: ECC + CONC group: 4.8 sets of 6 CONC repetitions and 6 ECC repetitions, CONC group: 4.8 sets of 12 CONC repetitions; intensity: ECC + CONC group: ECC and CONC MVC, CONC group: CONC MVC; tempo: 60°/s	12 weeks (3 sessions per week)	Increase in ECC KE strength (MVC) with ECC + CONC (36 %) and CONC (19 %), no change with CONC (-5 %), greater increase with ECC + CONC vs. CONC Increase in CONC KE strength (MVC) with ECC + CONC (25 %) and CONC (14 %), no change with CONC (-2 %), no difference between interventions Increase in lower body strength (back squat 3RM) with ECC + CONC (25 %) and CONC (15 %), no change with CONC (2 %), no difference between interventions Increase in lower body power (vertical jump; cm) with ECC + CONC (8 %), no change with CONC (3 %) or CONC (-1 %), no difference between interventions

Table 1 continued

Study, year	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Colliander and Tesch, [14] 1992	ECC + CONC $n = 10$, CONC $n = 8$, CONT $n = 7$ (25 M); mean age: 26.6 years; training status: untrained	KE (isokinetic; single-joint)	Volume: ECC + CONC group: 4.8 sets of 6 CONC repetitions and 6 ECC repetitions, CONC group: 4.8 sets of 12 CONC repetitions; intensity: ECC + CONC group: ECC and CONC MVC, CONC group: CONC MVC; tempo: 60°/s	12 weeks (3 sessions per week)	Increase in ECC KE strength (MVC) with ECC + CONC (37 %) and CONC (18 %), no change with CONT (NR), no difference between interventions Increase in CONC KE strength (MVC) with ECC + CONC (26 %) and CONC (13 %), no change with CONT (NR), no difference between interventions Increase in lower body strength (back squat 3RM) with ECC + CONC (23 %) and CONC (13 %), no change with CONT (NR), no difference between interventions Increase in lower body power (vertical jump; cm) with CONC (4 %), no change with ECC + CONC (8 %) or CONT (NR), no difference between interventions Increase in ECC KE strength (MVC; 60, 120 and 180 °/s) with ECC (29 %), no change with CONC (7 %) or CONT (-1 %) Increase in CONC KE strength (MVC; 180 °/s) with CONC (8 %), no change with ECC (1 %) or CONT (-5 %)
Duncan et al. [38], 1989	ECC $n = 16$, CONC $n = 14$, CONT $n = 18$ (48 M); mean age: 23.9 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 1 set of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 120°/s	6 weeks (2 sessions per week)	Increase in CONC ER strength (MVC; 60, 180, 210°/s) with ECC (NR) and CONC (NR), no difference between interventions Increase in CONC IR strength (MVC; 60, 180, 210°/s) with ECC (NR) and CONC (NR), no difference between interventions
Ellenbecker et al. [15], 1988	ECC $n = 11$, CONC $n = 11$ (22F and M); mean age: NR, university students; training status: varsity tennis athletes	ER and IR (isokinetic; single-joint)	Volume: 6 sets of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: pyramid across six sets (60, 180, 210, 180, 60°/s)	6 weeks (2 sessions per week)	Increase in ECC ER strength (MVC; 210°/s) with ECC (NR) and CONC (NR), no difference between interventions Increase in ECC IR strength (MVC; 60 and 180°/s) with CONC (8 %), no change with ECC (NR)
Elmer et al. [46], 2012	ECC $n = 6$, CONC $n = 6$ (12 M); mean age: 25.0 years; training status: untrained	HE and KE (isokinetic cycling; multi-joint)	Volume: 21 min; intensity: ECC group: 30 % concentric cycling peak power, CONC group: 19 % concentric cycling peak power; tempo: 60 rpm	7 weeks (3 sessions per week)	Increase in ECC ER strength (MVC; 210°/s) with ECC (NR) and CONC (NR), no difference between interventions Increase in ECC IR strength (MVC; 60 and 180°/s) with CONC (8 %), no change with ECC (NR) Increase in leg spring stiffness (kN/m) with ECC (10 %) vs. CONC (-2 %) Increase in jumping power (W) with ECC (7 %) vs. CONC (-2 %)
English et al. [16], 2014	ECC ₁₃₈ $n = 8$, ECC ₁₀₀ $n = 8$, CONC ₆₆ $n = 8$, CONC ₃₃ $n = 8$, CONC $n = 8$ (40 M); mean age: 34.9 years; training status: untrained	HE, KE and AE (isokinetic leg press; multi-joint)	Volume: 3.75 sets of 5 repetitions; intensity: ECC ₁₃₈ group: 138/76 % IRM, ECC ₁₀₀ group: 100/76 % IRM, CONC ₆₆ group: 66/76 % IRM, CONC ₃₃ : 33/76 % IRM, CONC: 0/76 % IRM; tempo: NR	8 weeks (3 sessions per week)	Increase in CONC HE and KE strength (leg press IRM) with ECC ₁₃₈ (20 %), ECC ₁₀₀ (13 %), CONC ₆₆ (8 %), CONC ₃₃ (8 %) and CONC (8 %), but ECC ₁₃₈ greater than CONC ₆₆ , CONC ₃₃ and CONC Increase in CONC AE strength (calf raise IRM) with ECC ₁₃₈ (11 %), ECC ₁₀₀ (12 %), CONC ₆₆ (7 %) and CONC ₃₃ (8 %), no change with CONC (5 %), no difference between interventions

Table 1 continued

Study, year	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Farthing and Chilitbeck [17], 2003	ECC _{Fast} , CONC _{Fast} $n = 13$ (9F, 4M), ECC _{Slow} , CONC _{Slow} $n = 11$ (4F, 7M), CONT $n = 10$ (8F, 2M); mean age: 22.2 years; training status: untrained	EF (isokinetic; single-joint)	Volume: 4.6 sets of 8 repetitions; intensity: ECC groups: ECC MVC, CONC groups: CONC MVC; tempo: fast groups: 180°/s, slow groups: 30°/s	8 weeks (3 sessions per week)	Increase in CONC EF strength (MVC) with ECC _{Fast} (23 %) greater than CONC _{Fast} (1 %), CONC _{Slow} (6 %) and CONT (0 %), but not ECC _{Slow} (16 %) Increase in ECC EF strength (MVC) with ECC _{Fast} (16 %) greater than CONC _{Fast} (-1 %), ECC _{Slow} (6 %), CONC _{Slow} (5 %) and CONT (0 %)
Farthing and Chilitbeck [44], 2003	ECC _{Fast} , CONC _{Fast} $n = 13$ (9F, 4M) ECC _{Slow} , CONC _{Slow} $n = 11$ (4F, 7M), CONT $n = 10$ (8F, 2M); mean age: 22.2 years; training status: untrained	EF (isokinetic; single-joint)	Volume: 4.6 sets of 8 repetitions; intensity: ECC groups: ECC MVC, CONC groups: CONC MVC; tempo: fast groups: 180°/s, slow groups: 30°/s	8 weeks (3 sessions per week)	Increase in contralateral ECC EF strength (MVC; 180 °/s) with ECC _{Fast} and CONC _{Fast} (23 %), no change with ECC _{Slow} , CONC _{Slow} (-17 %) or CONT (8 %), no difference between interventions
Farup et al. [18], 2014	ECC _{Whey} and ECC $n = 11$, CONC _{Whey} and CONC $n = 11$, within-subject design (22 M); mean age: 23.9 years; training status: untrained	KE (isoinertial; single-joint)	Volume: 9.3 sets of 10.7 repetitions; intensity: ECC groups: 90/75 % IRM, CONC groups: 75 % IRM; tempo: ECC: 2 s, CONC: 2 s	12 weeks (2.75 sessions per week)	Increase in CONC KE strength (MVC) with ECC (4 %), CONC _{Whey} (7 %) and CONC (20 %), no change with ECC _{Whey} (2 %), no difference between interventions Increase in ISO KE strength (MVC) with ECC _{Whey} (6 %) ECC (10 %), CONC _{Whey} (17 %) and CONC (20 %), no difference between interventions
Franchi et al. [37], 2014	ECC $n = 6$, CONC $n = 6$ (12 M); mean age: 25.0 years; training status: untrained	HE and KE (isokinetic; multi-joint)	Volume: 4 sets of 9 repetitions; intensity: ECC group: 80 % ECC IRM, CONC group: 80 % CONC IRM; tempo: ECC group: 3 s, CONC group: 2 s	10 weeks (3 sessions per week)	Increase in ECC KE strength (MVC) with ECC _{Whey} (10 %), ECC (8 %), CONC _{Whey} (8 %) and CONC (19 %), no difference between interventions Increase in ISO KE strength (MVC) with ECC (11 %) and CONC (9 %), no difference between interventions
Friedmann-Bette et al. [19], 2010	ECC $n = 14$, TRAD $n = 11$ (25 M); mean age: 24.4 years; training status: strength trained	KE (isoinertial; single-joint)	Volume: ECC group: 5 sets of 8 repetitions, TRAD group: 6 sets of 8 repetitions; intensity: ECC: 152/80 % IRM, TRAD: 80 % IRM; tempo: NR; explosive ECC and CONC	6 weeks (3 sessions per week)	Increase in CONC KE strength (IRM) with ECC (16 %) and TRAD (19 %), no difference between interventions Increase in squat jump (cm) with ECC (7 %), no change with TRAD (1 %)
Goddard et al. [20], 1998	ECC $n = 9$, TRAD $n = 9$, CONT $n = 10$ (17F, 21 M); mean age: 22.4 years; training status: recreationally active	KE (isokinetic; single-joint)	Volume: 1 set of 10 repetitions; intensity: ECC group: 120/80 % IRM, TRAD group: 80 % IRM; tempo: 30°/s	10 weeks (2)	Increase in CONC KE strength (MVC) with ECC (81 %) and TRAD (82 %) vs. CONT (7 %), no difference between interventions

Table 1 continued

Study, year	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Gross et al. [34], 2010	ECC $n = 8$, TRAD $n = 7$ (15 M); mean age: 17.6 years; training status: junior national skiers	HE and KE (isokinetic and isoinertial; multi-joint)	Volume: ECC group: 12 sets of 30 repetitions weight training and 20 min ECC cycling, TRAD group: 22.5 sets of 30 repetitions; intensity: ECC group: 40 % IRM weight training and 532 W ECC cycling, TRAD group: 40 % IRM; tempo: ECC cycling: 70 rpm, weight training: NR	6 weeks (3 sessions per week)	Increase in ISO HE and KE strength (leg press MVC) with ECC (10 %) and TRAD (12 %), no difference between interventions Decrease in squat jump (cm) with TRAD (-4 %), no change with ECC (2 %) Increase in countermovement jump (cm) with ECC (7 %), no change with TRAD (3 %)
Hawkins et al. [21], 1999	ECC $n = 8$, CONC $n = 8$ (within-subject design), CONT $n = 12$ (20F); mean age: 21.4 years; training status: untrained	KE and KF (isokinetic; single-joint)	Volume: ECC group: 3 sets of 3 repetitions, CONC group: 3 sets of 4 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: NR	18 weeks (3 sessions per week)	Increase in CONC KE strength (MVC) with ECC (18 %) and CONC (23 %), no change with CONT (-6 %), no difference between interventions Increase in ECC KE strength (MVC) with ECC (22 %) and CONC (17 %) increased, no change with CONT (-3 %), greater increase with ECC training vs. CONC Increase in CONC KF strength (MVC) ECC (13 %), no change with CONC (6 %) or CONT (NR) Increase in ECC KF strength (MVC) with ECC (14 %) and CONC (13 %), no change with CONT (NR), no difference between interventions
Higbie et al. [39], 1996	ECC $n = 19$, CONC $n = 16$, CONT $n = 19$ (54F); mean age: 20.5 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 3 sets of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 60°/s	10 weeks (3 sessions per week)	Increase in CONC KE strength (MVC) with CONC (18 %), no change with ECC (7 %) or CONT (5 %), greater increase with CONC training vs. ECC Increase in ECC KE strength (MVC) with ECC (36 %) and CONC (13 %), no change with CONT (-2 %), greater increase with ECC training vs. CONC Increase in CONC KE activation (EMG) with CONC (22 %) greater than CONT (-8 %), no change with ECC (7 %) Increase in ECC KE activation (EMG) with ECC (17 %) and CONC (20 %) greater than CONT (-9 %), no difference between interventions

Table 1 continued

Study, year	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Hortobagyi et al. [22], 1996	ECC $n = 7$, CONC $n = 8$, CONT $n = 6$ (21 M); mean age: 21.3 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 5.3 sets of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 60°/s	12 weeks (3 sessions per week)	Increase in CONC KE strength (MVC) with ECC (33 %) and CONC (53 %), no change with CONT (NR), no difference between interventions Increase in ECC KE strength (MVC) with ECC (116 %), no change with CONC (40 %) or CONT (NR), greater increase with ECC training vs. CONC Increase in CONC KE activation (EMG) with CONC (28 %), no change with ECC (62 %) or CONT (NR), no difference between interventions Increase in ECC KE activation (EMG) with ECC (188 %), no change with CONC (10 %) or CONT (NR), greater increase with ECC training vs. CONC
Hortobagyi et al. [45], 1997	ECC $n = 7$, CONC $n = 8$, CONT $n = 6$ (21 M); mean age: 21.3 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 5.3 sets of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 60°/s	12 weeks (3 sessions per week)	Increase in contralateral CONC KE Strength (MVC) with CONC (30 %), no change with ECC (18 %) or CONT (0 %), no difference between interventions Increase in contralateral ISO KE strength (MVC) with ECC (39 %) and CONC (22 %), no change with CONT (2 %), greater increase with ECC vs. CONC Increase in contralateral ECC KE strength (MVC) with ECC (77 %), no change with CONC (18 %) or CONT (4 %), greater increase with ECC vs. CONC
Hortobagyi et al. [23], 2000	ECC $n = 12$, ECC + CONC $n = 12$, CONC $n = 12$, CONT $n = 24$ (24F, 24 M); mean age: 22.0 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 5.3 sets of 10 repetitions; intensity: ECC group: ECC MVC, ECC + CONC group: ECC and CONC MVC, CONC group: CONC MVC; tempo: 60°/s	12 weeks (3 sessions per week)	Increase in CONC KE strength (MVC) with ECC (25 %), ECC + CONC (40 %) and CONC (44 %), no change with CONT (NR), no difference between interventions Increase in ISO KE strength (MVC) with ECC (42 %), ECC + CONC (38 %) and CONC (31 %), no change with CONT (NR), greater increase with ECC and ECC + CONC vs. CONC Increase in ECC KE strength (MVC) with ECC (86 %), ECC + CONC (70 %) and CONC (20 %), no change with CONT (NR), no difference between interventions
Kaminski et al. [24], 1998	ECC $n = 9$, CONC $n = 9$, CONT $n = 9$ (27 M); mean age: 22.9 years; training status: untrained	KF (isoinertial; single-joint)	Volume: 2 sets of 8 repetitions; intensity: ECC group: 100/40 % 1RM, CONC group: 80 % 1RM; tempo: NR	6 weeks (2 sessions per week)	Increase in CONC KF strength (1RM/BW) with ECC (29 %) and CONC (19 %), no change with CONT (5 %), no difference between interventions Increase in ECC KF strength (MVC) with ECC (30 %), no change with CONC (13 %) or CONT (-5 %), no difference between interventions

Table 1 continued

Study, year	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Komi and Buskirk [25], 1972	ECC $n = 11$, CONC $n = 10$, CONT $n = 10$ (31 M); mean age: 19.6 years; training status: untrained	EF (isokinetic; single-joint)	Volume: 1 set of 6 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: NR	7 weeks (4 sessions per week)	Increase in CONC EF strength (MVC) with ECC (16 %) and CONC (12 %) vs. CONT (2 %), no difference between interventions Increase in ISO EF strength (MVC) with ECC (7 %) vs. CONT (1 %), no change with CONC (6 %) Increase in ECC EF strength (MVC) with ECC (16 %) and CONC (7 %) vs. CONT (-4 %), greater increase with ECC vs. CONC
LaStayo et al. [35], 1999	ECC $n = 4$, CONC $n = 5$ (5F, 4 M); mean age: 21.5 years; training status: untrained	HE and KE (isokinetic cycling; multi-joint)	Volume: 27.5 min; intensity: ECC group: 1 L/min O ₂ consumption, CONC group: 1.2 L/min O ₂ consumption; tempo: 55 rpm	6 weeks (4.2 sessions per week)	Increase in ISO KE strength (MVC) with ECC (27 %), no change with CONC (10 %)
LaStayo et al. [36], 2000	ECC $n = 7$, CONC $n = 7$ (14 M); mean age: 23.9 years; training status: untrained	HE and KE (isokinetic cycling; multi-joint)	Volume: 27.5 min; intensity: 62 % maximum heart rate; tempo: 60 rpm	8 weeks (3.5 sessions per week)	Increase in ISO KE strength (MVC) with ECC (36 %), no change with CONC (2 %)
Liu et al. [47], 2013	ECC + CONC _{Fast} $n = 10$, ECC + CONC _{Slow} $n = 10$, TRAD $n = 10$ (30 M); mean age: 19.5 years; training status: untrained	HE and KE (ECC groups: isokinetic, TRAD group: isoinertial; multi-joint)	Volume: 5 sets of 10 repetitions; intensity: ECC groups: ECC MVC, TRAD group: 70 % 1RM; tempo: ECC _{Fast} : 2.5 Hz, ECC _{Slow} and TRAD: 0.5 Hz	10 weeks (3 sessions per week)	Increase in lower body power (vertical jump; cm) with ECC + CONC _{Fast} (4 %) and ECC + CONC _{Slow} (3 %), no change with TRAD (2 %), greater increase with ECC + CONC _{Fast} vs. ECC + CONC _{Slow} and TRAD Increase in drop jump (cm) with ECC + CONC _{Fast} (6 %), no change with ECC + CONC _{Slow} (1 %) or TRAD (1 %), greater increase with ECC + CONC _{Fast} vs. ECC + CONC _{Slow}
Malliaras et al. [26], 2013	ECC _{Heavy} $n = 10$, ECC _{Light} $n = 10$, CONC $n = 9$, CONT $n = 9$ (38 M); mean age: 27.5 years; training status: untrained	KE (isoinertial; single-joint)	Volume: ECC _{Heavy} and CONC groups: 4 sets of 7.5 repetitions, ECC _{Light} group: 4 sets of 13.5 repetitions; intensity: ECC _{Heavy} group: 80 % ECC 1RM, ECC _{Light} and CONC groups: 80 % CONC 1RM; tempo: ECC: 5 s, CONC: 1 s	12 weeks (3 sessions per week)	Decrease in 30 m sprint time (s) with ECC + CONC _{Fast} (-0.23 %), ECC + CONC _{Slow} (-0.12 %) and TRAD (-0.12 %), greater improvement with ECC + CONC _{Fast} vs. ECC + CONC _{Slow} and TRAD Increase in stretch shortening cycle efficiency with ECC + CONC _{Fast} (11 %), no change with ECC + CONC _{Slow} (4 %) or TRAD (2 %), greater increase with ECC + CONC _{Fast} vs. ECC + CONC _{Slow} and TRAD Increase in CONC KE strength (5RM) with ECC _{Heavy} (77 %), ECC _{Light} (61 %) and CONC (53 %), no change with CONT (NR), greater increase with ECC _{Heavy} vs. ECC _{Light} and CONC

Table 1 continued

Study, year	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Miller et al. [27], 2006	ECC $n = 17$, CONC $n = 21$ (38F); mean age: 20.0 years; training status: untrained	KF and KE (isokinetic; single-joint)	Volume: 4.5 sets of 6 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 60°/s	20 weeks (3 sessions per week)	Increase in CONC KE strength (MVC) with ECC (15 %) and CONC (10 %), no difference between interventions Increase in CONC KF strength (MVC) with ECC (29 %) and CONC (27 %), no difference between interventions Increase in ECC KE strength (MVC) with ECC (28 %) and CONC (12 %), greater increase with ECC vs. CONC Increase in ECC KF strength (MVC) with ECC (40 %) and CONC (20 %), greater increase with ECC vs. CONC Increase in CONC KE RFD (ms) with ECC (14 %) and CONC (14 %), no difference between interventions Increase in CONC KF RFD (ms) with ECC (35 %) and CONC (21 %), no difference between interventions Increase in ECC KE RFD (ms) with ECC (26 %) and CONC (2 %), greater increase with ECC vs. CONC Increase in ECC KF RFD (ms) with ECC (21 %) and CONC (9 %), greater increase with ECC vs. CONC Increase in CONC ER strength (MVC) with ECC (9 %) and CONC (7 %) vs. CONC (-9 %), no difference between interventions
Mont et al. [28], 1994	ECC $n = 8$, CONC $n = 9$, CONT $n = 13$ (30 M); mean age: 33.0 years; training status: elite tennis players	ER and IR (isokinetic; single-joint)	Volume: 8 sets of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: pyramid across eight sets (90, 120, 150, 180, 180, 180, 120, 90°/s)	6 weeks (3 sessions per week)	Increase in CONC IR strength (MVC) with ECC (2 %) and CONC (12 %) vs. CONC (-9 %), no difference between interventions Increase in ECC ER strength (MVC) with ECC (18 %) and CONC (10 %) vs. CONC (-1 %), no difference between interventions Increase in ECC IR strength (MVC) with ECC (5 %) and CONC (18 %) vs. CONC (-4 %), no difference between interventions
Moore et al. [29], 2012	ECC and CONC $n = 9$, within-subject design (9 M); mean age: 22.0 years; training status: untrained	EF (isokinetic; single-joint)	Volume: ECC group 4.44 sets of 10 repetitions, CONC group: 4.44 sets of 14 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 45°/s	9 weeks (2 sessions per week)	Increase in CONC EF strength (MVC) with ECC (11 %) and CONC (14 %), no difference between interventions Increase in ISO EF strength (MVC) with ECC (9 %) and CONC (19 %), no difference between interventions Increase on ECC EF strength (MVC) with ECC (8 %) and CONC (11 %), no difference between interventions

Table 1 continued

Study, year	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Nickols-Richardson et al. [30], 2007	ECC $n = 33$, CONC $n = 37$ (70F); mean age: 35.0 years; training status: untrained	KF, KE, EF and EE (isokinetic; single-joint)	Volume: 4.5 sets of 6 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 60°/s	20 weeks (3 sessions per week)	Increase in CONC KF and KE strength (MVC) with ECC (29 %) and CONC (19 %), no difference between interventions Increase in CONC EF and EE strength (MVC) with ECC (25 %) and CONC (13 %), greater increase with ECC vs. CONC Increase in CONC KE strength (MVC; 30 and 90°/s) with CONC (12 %), no change with ECC (4 %) Increase in ISO KE strength (MVC) with CONC (14 %), no change with ECC (6 %) Increase in ECC KE strength (MVC; 30 and 90°/s) with ECC (24 %) and CONC (13 %), no difference between interventions Increase in ECC KE strength (MVC) with ECC (26 %) and CONC (23 %), no difference between interventions
Seger et al. [40], 1998	ECC $n = 5$, CONC $n = 5$ (10 M); mean age: 24.5 years; training status: moderately trained	KE (isokinetic; single-joint)	Volume: 4 sets of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 90°/s	10 weeks (3 sessions per week)	Increase in CONC KE strength (MVC) with CONC (14 %), no change with ECC (6 %) Increase in ECC KE strength (MVC; 30 and 90°/s) with ECC (24 %) and CONC (13 %), no difference between interventions
Spurway [42], 2000	ECC and CONC $n = 20$, within-subject design (10F, 10 M); mean age: 24.0 years; training status: untrained	KE (isoinertial; single-joint)	Volume: 3 sets of 6 repetitions; intensity: ECC group: 85 % ECC IRM, CONC group: 85 % CONC IRM; tempo: NR	6 weeks (3 sessions per week)	Increase in ECC KE strength (MVC) with CONC (26 %) and CONC (23 %), no difference between interventions
Tomberlin et al. [41], 1991	ECC $n = 21$, CONC $n = 19$, CONT $n = 23$ (32F, 31 M); mean age: 27.1 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 3 sets of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 100°/s	6 weeks (3 sessions per week)	Increase in CONC KE strength (MVC) with CONC (8 %), no change with ECC (4 %) or CONT (-2 %), greater increase with CONC vs. ECC and CONT Increase in ECC KE strength (MVC) with ECC (21 %) and CONC (11 %), no change with CONT (5 %), greater increase with ECC vs. CONC and CONT Increase in CONC EF strength (IRM) with ECC (14 %) and CONC (18 %), no difference between interventions Increase in ECC EF strength (IRM) with ECC (9 %) and CONC (3 %), greater increase with ECC vs. CONC
Vikne et al. [31], 2006	ECC $n = 9$, CONC $n = 8$ (17 M); mean age: 27.1 years; training status: resistance trained	EF (isoinertial; single-joint)	Volume: 3.9 sets of 6 repetitions; intensity: ECC group: 94 % ECC IRM, CONC group: 94 % CONC IRM; tempo: ECC: 3.5 s, CONC: explosive	12 weeks (2.5 sessions per week)	Increase in lower body strength (back squat IRM) with ECC (19 %) and CONC (25 %), no difference between interventions Increase in upper body strength (bench press IRM) with ECC (9 %) and CONC (10 %), no difference between interventions
Yarrow et al. [32], 2008	ECC $n = 10$, CONC $n = 12$ (22 M); mean age: 22.1 years; training status: untrained	Lower Body/Back Squat and Upper Body/Bench Press (isoinertial; multi-joint)	Volume: ECC group: 3 sets of 6 repetitions, CONC group: 4 sets of 6 repetitions; intensity: ECC group: 110/44 % IRM, CONC group: 65 % IRM; tempo: 6 s per repetition	5 weeks (3 sessions per week)	Increase in lower body strength (back squat IRM) with ECC (19 %) and CONC (25 %), no difference between interventions Increase in upper body strength (bench press IRM) with ECC (9 %) and CONC (10 %), no difference between interventions

%/s degrees per second, *cm* centimetres, *AE* ankle extensors, *CONC* concentric, *CONT* control, *ECC* eccentric, *ECC + CONC* coupled maximal eccentric and concentric contractions, *EE* elbow extensors, *EF* elbow flexors, *EMG* electromyography, *ER* shoulder external rotators, *F* females, *HE* hip extensors, *IR* shoulder internal rotators, *ISO* isometric, *KE* knee extensors, *KF* knee flexors, *kN/m* kilonewtons per metre, *L/m* litres per minute, *M* males, *ms* milliseconds, *MVC* maximum voluntary contraction, *n* number of subjects, $N \cdot s^{-1}$ newtons per second, *NR* not reported, *RFD* rate of force development, *RM* repetition maximum, *rpm* revolutions per minute, *s* seconds, *TRAD* traditional resistance training, *W* watts

performance and leg spring stiffness have been found to increase following eccentric, but not concentric or traditional, resistance training [46, 47]. Squat jump performance, which involves no SSC component and therefore reflects muscle power independent of elastic energy utilization and stretch reflex potentiation, may also benefit to a greater extent following eccentric training compared with concentric or traditional resistance training [19, 34]. Aligning with observations on muscle strength, fast eccentric training appears to have a more potent effect on measures of muscle power than slow eccentric training. Vertical jump, drop jump, SSC efficiency and sprinting performance (i.e. 30 m) all improved to a greater extent following fast (i.e. 2.5 Hz isokinetic squat movement) combined eccentric/concentric training compared with slow combined eccentric/concentric (i.e. 0.5 Hz isokinetic squat movement) and traditional resistance training [47]. Improvements in contractile RFD appear to be dependent on the method of assessment. Isometric RFD from 0 to 30 ms increased to a greater extent with concentric versus eccentric training [33], while eccentric training has been found to elicit a greater increase in eccentric RFD (i.e. as inferred from time to peak torque during isokinetic knee extension and flexion) and an equivalent increase in concentric RFD compared with concentric training [27].

3.2.1 Muscle Strength

The mechanisms underpinning strength improvements with eccentric training are likely a combination of neural, morphological and architectural factors (Sect. 3.3), although relatively few studies have investigated the possible neural adaptations following eccentric training [22, 25, 39]. The mode-specificity of strength improvements in conjunction with cross-education supports the importance of neural mechanisms, while other evidence indicates that changes in eccentric strength are attained via increased agonist voluntary activation [48, 49] and decreased antagonist coactivation [50]. The inability to fully activate muscle during eccentric contractions [51], particularly in untrained subjects [52], may explain the large improvements in eccentric strength and the observation that eccentric training increases eccentric strength to a greater degree than concentric training increases concentric strength [22, 39]. In support of an improved neural drive, Vangsgaard et al. [49] found 5 weeks of eccentric training of the trapezius muscle increased muscle excitability (i.e. inferred from maximal evoked H-reflex) in concert with a 26 % increase in MVC, aligning with previous findings [53]. Improvements in agonist voluntary activation during eccentric contractions may result from a disinhibition of presynaptic Golgi Ib and joint afferents known to inhibit excitatory muscle spindle Ia afferents [54]. Surface

electromyography (EMG) has been found to increase during eccentric contractions, but not concentric contractions, following eccentric training [22, 39], although conflicting findings have been reported [39]. Maximal EMG (i.e. peak root mean square [RMS] values and integrated voltage from the EMG [iEMG]) following training is proposed to reflect the electrical excitation of muscle, which is influenced by the number and size (i.e. type I vs. type II fibres) of recruited motor units, motor unit discharge rate and synchrony [22, 39]. Based on findings using the twitch interpolation technique, motor unit activation during an isometric MVC can be, although not in all cases, maximal in untrained subjects during single joint movements [55]. It has been proposed that increases in EMG with eccentric training result from increases in motor unit discharge rate [39]. Contrary to reports on isometric contractions, maximal activation is typically inhibited during maximal eccentric contractions with untrained subjects [54], which indicates that both motor unit recruitment and discharge rates could contribute to increases in strength. The relative contribution of motor unit activation to the voluntary activation deficit may increase during compound multi-joint movements [55], but methodological limitations with measuring motor unit activity make it challenging to corroborate this supposition. Nonetheless, it is believed that the motor unit discharge rate is the primary inhibitory mechanism constraining voluntary activation during eccentric contractions [52], and thus the neural factor that contributes most to increased eccentric strength following eccentric, or heavy, resistance training [54].

3.2.2 Muscle Power and Stretch-Shortening Cycle Performance

Increases in muscle power following chronic resistance training are generally expected, particularly for subjects with a limited training history [56]. Muscle power is dictated by the force–velocity relationship and therefore the capacity of muscle to both produce force and shorten rapidly [3]. The neural mechanisms underpinning improvements in strength, along with morphological and architectural adaptations (Sect. 3.3), probably contribute to improvements in muscle power with eccentric training [3]. Increases in muscle power will largely arise from improvements in the capability to rapidly recruit larger motor units (i.e. type IIa and IIx) and increases in motor unit firing frequency [3]. Faster recruitment of larger motor units may be achieved by lowering the threshold of recruitment [57], or via a preferential recruitment. Experimental evidence supporting a preferential recruitment is not compelling [52], and therefore a lower threshold of recruitment seems most plausible in explaining a training-induced improvement in rapid motor unit recruitment [3].

Increases in motor unit firing frequency could also contribute to enhancements in eccentric power (i.e. during a vertical jump), and would align with the purported effects of motor unit discharge rates on eccentric strength [54]. Improvements in motor unit synchronization (i.e. intramuscular coordination) and intermuscular coordination (i.e. improved synergist coactivation and decreased antagonist coactivation) could also play a role in the expression of neuromuscular power [3, 58]. Improved eccentric force control is a postulated adaptation to eccentric training [56] and it is possible that increased eccentric coordination could contribute to SSC performance. SSC performance during various outcome measures (i.e. vertical jump, drop jump and SSC efficiency) can be enhanced by modulating the eccentric phase of the movement [46, 47, 59, 60]. Papadopoulos and colleagues [59] found an eccentric leg-press training protocol elicited smaller changes in ankle, hip and knee angles during the eccentric phase of a drop jump that corresponded to improved jump height, power and contact time. The increased joint stiffness during the eccentric phase of the SSC probably enhanced the utilization of elastic energy [59] and allowed the muscle to operate closer to its optimum length and shortening velocity [60]. These adaptations appear to translate to functional performance such as sprinting, particularly with inclusion of fast mixed eccentric/concentric contractions in the training protocol [47]. Cook et al. [61] found that 3 weeks of controlled tempo eccentric training in semi-professional rugby players improved back squat, bench press and countermovement jump performance to a greater extent than traditional resistance training. However, the authors found that the addition of overspeed sprint and power training was necessary to elicit increases in 40-m sprint performance with eccentric training [61]. A 3.3 % increase in flying 10 m sprint time has also been reported in well-trained soccer players following 10 weeks of eccentric flywheel training [62], but contraction speed was not reported. While further research is necessary, it may be postulated that fast contraction velocities optimize the transfer of eccentric-related adaptations to sprint performance involving a fast SSC component.

3.2.3 Contractile Rate of Force Development

The lack of studies investigating the effects of eccentric training on RFD, combined with divergent methods of assessment, make it difficult to draw firm conclusions in this area. Contractile RFD is influenced by muscle strength, cross-sectional area (CSA), fibre type and myosin heavy chain (MHC) composition, MTU viscoelastic properties, and neural drive [63, 64]. The relative contribution of these factors probably depends on the RFD time interval assessed. A combination of neural drive and intrinsic muscle

properties (i.e. MHC composition) appears to play a marked role during early-phase (i.e. <100 ms) RFD [65], which is unsurprising given the cross-bridge cycling rates of both IIX and IIA fibres are substantially greater than type I fibres [66, 67]. Improvements in late-phase (i.e. >100 ms) RFD may be more related to neural drive, muscle CSA and tendon/aponeurosis stiffness [65]. Indeed, Andersen and Aagaard [64] found that very early RFD (i.e. <40 ms) was related to twitch RFD more so than maximal strength, and from 90 ms onwards strength accounted for 52–81 % of the variance in voluntary RFD. Furthermore, tendon and aponeurosis stiffness has been demonstrated to account for up to 30 % of the variance in RFD from 0 to 200 ms [63]. Training interventions that change these underlying mechanisms could plausibly influence early- and late-phase RFD to differing extents [68]. Six weeks of fast concentric (i.e. 180°/s) training can improve early-phase RFD by 33–56 %, independent of any change in late-phase RFD or strength, a finding that has been attributed to changes in neural drive [69]. Fast eccentric (i.e. 180°/s) resistance training has been similarly demonstrated to improve early-phase RFD by 30 %, but also in conjunction with a 28 % improvement in strength [65]. The mechanisms by which eccentric training improve early-phase RFD may be a combination of intrinsic muscle and neural adaptations. Similar to improvements in muscle strength and power, the unique neural strategies involved with eccentric training may improve RFD via both increased volitional supraspinal drive and spinal reflex disinhibition [65]. Late-phase improvements in RFD have been demonstrated to occur in parallel with improvements in strength with heavy resistance training [64], and therefore the increases in strength observed with eccentric training may contribute to enhanced late-phase RFD. Isometric training that includes the intention to contract explosively (i.e. ‘ballistic-intended’) has been found to improve both early and late RFD [70], and it has been proposed that the intention to act explosively is more important than the type of contraction performed [69]. Further research is necessary to determine best practice in maximizing improvements in RFD; as long as there is maximal intent to contract explosively, eccentric, isometric and concentric training protocols can lead to performance enhancements.

3.2.4 Summary of Eccentric Training Effects on Muscle Mechanical Function

In summary, eccentric training may improve overall strength to a greater extent than concentric and traditional modalities, although there is a mode-specificity (i.e. contraction type and velocity) of improvements [8]. Increases in strength result from a combination of neural, morphological and architectural adaptations. Increased voluntary

activation of agonists during eccentric contractions via a disinhibition of excitatory inflow to spinal motor neurons may underpin the marked increases in eccentric strength observed following training [54]. Based on EMG experiments, an increase in motor unit discharge rate appears to play a more important role than changes in motor unit recruitment [39], although further research is necessary to determine whether this observation holds during compound, multi-joint movements [55]. Eccentric training improves muscle power and SSC performance to a greater extent than concentric or traditional modalities, while reports on changes in RFD have been mixed. Improvements are likely related to increases in eccentric and concentric strength, an improved ability to rapidly recruit fast motor units, a shift towards a faster muscle phenotype (Sect. 3.3), and an enhanced eccentric phase within the SSC. Given the mode-specificity of adaptations, dedicated concentric training should be performed in addition to eccentric training if improvements in concentric strength, and possibly power, are of importance. In practice, traditional resistance training combining eccentric and concentric actions remains most prevalent. The use of traditional exercises with additional overload above the concentric maximum during the eccentric phase may therefore maximize outcomes across concentric, isometric and eccentric strength [16]. Furthermore, fast eccentric velocities may induce greater strength, power and SSC improvements than slow eccentric velocities [17, 47]. It has also been suggested that the optimal duration between the cessation of eccentric training and performance assessment may necessarily differ from concentric training in expressing improvements in muscle mechanical function [71]. More research is needed to determine the fatigue, recovery and adaptation time-course, as well as subsequent performance responses, with eccentric training.

3.3 Muscle-Tendon Unit Morphology and Architecture

Eccentric [11, 13, 16, 17, 19–21, 29, 31, 34, 36, 37, 39, 40, 72, 73], concentric [11, 13, 17, 29, 30, 37, 72, 73] and traditional [19, 20, 39] resistance training have all been found to increase muscle size using a variety of measures (i.e. dual-energy x-ray absorptiometry [DXA], muscle circumference, magnetic resonance imaging [MRI], peripheral computerized tomography [pQCT] and ultrasound) (Table 2). Muscle hypertrophy is a common finding with eccentric and concentric training modalities. Either a greater increase [16, 17, 21, 34, 36, 39, 40], or no difference [11, 13, 29, 30, 37, 72, 73] has been reported following

eccentric training. Those comparing eccentric training with traditional resistance training reported no differences between modalities [19, 20]. Contraction type appears to mediate a region-specific hypertrophy; eccentric training tends to induce greater increases in distal muscle size, while mid-muscle hypertrophy occurs to a greater extent following concentric training [37, 40]. It is plausible that reported literature is not entirely representative of the extent of adaptation with eccentric training, given that muscle morphology and architecture are typically assessed at the mid-belly of the muscle. When looking at fibre-type-specific responses (i.e. via muscle biopsy and electron microscopy), greater increases in type II fibre area have been observed with eccentric versus concentric or traditional training [19, 22, 23, 31]. In particular, an improved maintenance of, or increase in, type IIX fibre area has been reported with eccentric training relative to concentric [13] or traditional resistance training [19, 23]. Fibre type composition may be uniquely influenced by eccentric training, and either an improved maintenance of [13, 31], or similar reduction of [22], IIX fibres has been found compared with concentric training. As per changes in muscle strength, the intensity and velocity of muscle contraction influence the magnitude of muscle hypertrophy with eccentric training. Even when all conditions involve supra-maximal loads (i.e. greater than maximal concentric strength), greater increases in hypertrophy have been found with heavier eccentric training conditions [16], while fast eccentric velocities (i.e. 180°/s) also induced larger increases in muscle CSA [17]. The addition of whey protein hydrolysate supplementation in the post-exercise period has been found to increase [73], or have no effect on [72], the hypertrophic response following eccentric training. Few studies investigated changes in muscle architecture [11, 37] or tendon structure and function [26, 72] with differing contraction types. Vastus lateralis fascicle length was found to increase with both eccentric and concentric training [11], or eccentric training only [37]. Conflicting results were reported for vastus lateralis fascicle angle, with increases observed following eccentric training only [11], or concentric training only [37]. The patellar tendon cross-sectional area can increase with eccentric training, while the addition of whey protein hydrolysate supplementation appears necessary to promote increases with concentric training [72]. Patellar tendon Young's modulus at 50–75 % MVC has been demonstrated to increase with both eccentric and concentric training, but only with eccentric training at 75–100 % MVC [26]. Maximal tendon force and stress also increased with eccentric, but not concentric, training. Furthermore, a particularly marked increase was observed with heavier eccentric loading [26].

Table 2 Studies comparing the effects of eccentric-overload and coupled maximal eccentric and concentric contractions with traditional and concentric-only resistance training on muscle tendon unit morphology and architecture

Study	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Blazevich et al. [11]	ECC $n = 11$, CONC $n = 10$, CONT $n = 9$ (16F, 14 M); mean age: 22.8 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 5 sets of 6 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 30°/s	10 weeks (3 sessions per week)	Increase in quadriceps CSA (cm^3) with ECC and CONC (average change: 10 %), no change with CONT (NR), no difference between conditions Increase in quadriceps fascicle length (mm) with ECC (3 %) and CONC (6 %), no change with CONT (NR), no difference between conditions Increase in vastuslateralis fascicle angle ($^\circ$) with ECC (21 %), no change with CONC (13 %) or CONT (NR), no difference between interventions
Collander and Tesch [13]	ECC + CONC $n = 11$, CONC $n = 11$, CONT $n = 7$ (29 M); mean age: 26.3 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 4.8 sets of 12 repetitions; intensity: ECC + CONC group: ECC and CONC MVC, CONC group: CONC MVC; tempo: 60°/s	12 weeks (3 sessions per week)	Increase in thigh CSA (mm) with ECC + CONC (1 %) and CONC (1 %), no change with CONT (0.2 %), no difference between interventions Decrease in quadriceps type IIx area (μm^2) with CONC (-30 %), no change with ECC + CONC (-34 %) or CONT (-33 %), no difference between interventions
English [16]	ECC ₁₃₈ $n = 8$, ECC ₁₀₀ $n = 8$, CONC ₆₆ $n = 8$, CONC ₃₃ $n = 8$, CONC $n = 8$ (40 M); mean age: 34.9 years; training status: untrained	HE, KE and AE (isokinetic leg press; multi-joint)	Volume: 3.75 sets of 5 repetitions; intensity: ECC ₁₃₈ group: 138/76 % IRM, ECC ₁₀₀ group: 100/76 % IRM, CONC ₆₆ group: 66/76 % IRM, CONC ₃₃ : 33/76 % IRM, CONC: 0/76 % IRM; tempo: NR	8 weeks (3 sessions per week)	Decrease in quadriceps type IIx fibre composition (%) with CONC (-43 %), no change with ECC + CONC (-49 %) or CONT (-75 %), no difference between interventions Increase in leg CSA (kg) with ECC ₁₃₈ (2.4 %), no change with ECC ₁₀₀ (1.5 %), CONC ₆₆ (2.2 %), CONC ₃₃ (1.5 %) or CONC (-1.5 %), no difference between conditions
Farthing and Chilibeck [17]	ECC _{Fast} , CONC _{Fast} $n = 13$ (9F, 4 M) ECC _{Slow} , CONC _{Slow} $n = 11$ (4F, 7 M), CONT $n = 10$ (8F, 2 M); mean age: 22.2 years; training status: untrained	EF (isokinetic; single-joint)	Volume: 4.6 sets of 8 repetitions; intensity: ECC groups: ECC MVC, CONC groups: CONC MVC; tempo: Fast groups: 180°/s, Slow groups: 30°/s	8 weeks (3 sessions per week)	Increase in biceps CSA (mm) with ECC _{Fast} (13 %), CONC _{Fast} (3 %), ECC _{Slow} (8 %), CONC _{Slow} (5 %), no change with CONT (-1 %), greater increase with ECC _{Fast} vs. CONC _{Fast} , CONC _{Slow} and CONT
Farup et al. [18]	ECC _{Whey} and ECC $n = 11$, CONC _{Whey} and CONC $n = 11$, within-subject design (22 M); mean age: 23.9 years; training status: untrained	KE (isoinertial; single-joint)	Volume: 9.3 sets of 10.7 repetitions; intensity: ECC groups: 90/75 % IRM, CONC groups: 75 % IRM; tempo: ECC: 2 s, CONC: 2 s	12 weeks (2.75 sessions per week)	Increase in quadriceps type I fibre CSA (μm^2) with ECC _{Whey} (14 %), ECC (16 %), CONC _{Whey} (22 %) and CONC (12 %), no difference between interventions Increase in quadriceps type II fibre CSA (μm^2) with CONC _{Whey} (25 %), no change for ECC _{Whey} (1 %), ECC (11 %) or CONC (7 %), greater increase with CONC _{Whey} vs. ECC _{Whey} and CONC

Table 2 continued

Study	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Farup et al. [72]	ECC _{whey} and ECC $n = 11$, CONC _{whey} and CONC $n = 11$, within-subject design (22 M); mean age: 23.9 years; training status: untrained	KE (isoinertial; single-joint)	Volume: 9.3 sets of 10.7 repetitions; intensity: ECC groups: 90/75 % IRM, CONC groups: 75 % IRM; tempo: ECC: 2 s, CONC: 2 s	12 weeks (2.75 sessions per week)	Increase in quadriceps CSA (cm ²) with ECC _{whey} (6 %), ECC (2 %), CONC _{whey} (5 %) and CONC (3 %), no difference between interventions Increase in patellar tendon CSA (cm ²) with ECC _{whey} (15 %), ECC (10 %) and CONC _{whey} (15 %), no change with CONC (7 %), no difference between interventions
Franchi et al. [37]	ECC $n = 6$, CONC $n = 6$ (12 M); mean age: 25.0 years; training status: untrained	HE and KE (isokinetic; multi-joint)	Volume: 4 sets of 9 repetitions; intensity: ECC group: 80 % ECC IRM, CONC group: 80 % CONC IRM; tempo: ECC group: 3 s, CONC group: 2 s	10 weeks (3 sessions per week)	Increase in quadriceps CSA (cm ³) with ECC (6 %) and CONC (8 %), no difference between interventions
Friedmann-Bette et al. [19]	ECC $n = 14$, TRAD $n = 11$ (25 M); mean age: 24.4 years; training status: strength trained	KE (isoinertial; single-joint)	Volume: ECC group: 5 sets of 8 repetitions, TRAD group: 6 sets of 8 repetitions; intensity: ECC: 152/80 % IRM, TRAD: 80 % IRM; tempo: NR; explosive ECC and CONC	6 weeks (3 sessions per week)	Increase in quadriceps fascicle length (cm) with ECC (12 %) and CONC (5 %), greater increase with ECC vs. CONC Increase in vastuslateralis fascicle angle (°) with CONC (30 %), no change with ECC (5 %), greater increase with CONC vs. ECC Increase in quadriceps CSA (cm ²) with ECC (6 %) and TRAD (8 %), no difference between interventions Increase in quadriceps type Ix fibre CSA (µm ²) with ECC (20 %), no change with TRAD (10 %), no difference between interventions
Godard et al. [20]	ECC $n = 9$, TRAD $n = 9$, CONC $n = 10$ (12F, 16 M); mean age: 22.4 years; training status: recreationally active	KE (isokinetic; single-joint)	Volume: 1 set of 10 repetitions; intensity: ECC group: 120/80 % IRM, TRAD group: 80 % IRM; tempo: 30°/s	10 weeks (2 sessions per week)	Increase in thigh CSA (mm) with ECC (5 %) and TRAD (6 %) vs. CONC (1 %), no difference between interventions
Gross et al. [34]	ECC $n = 8$, TRAD $n = 7$ (15 M); mean age: 17.6 years; training status: junior national skiers	HE and KE (isokinetic and isoinertial; multi-joint)	Volume: ECC group: 12 sets of 30 repetitions weight training and 20 min ECC cycling, TRAD group: 22.5 sets of 30 repetitions; intensity: ECC group: 40 % IRM weight training and 532 W ECC cycling, TRAD group: 40 % IRM; tempo: ECC cycling: 70 rpm, weight training: NR	6 weeks (3 sessions per week)	Increase in leg lean mass (g) with ECC (2 %), no change with TRAD (NR), no difference between interventions
Hawkins et al. [21]	ECC $n = 8$, CONC $n = 8$ (within-subject design), CONC $n = 12$ (20F); mean age: 21.4 years; training status: untrained	KF and KE (isokinetic; single-joint)	Volume: ECC group: 3 sets of 3 repetitions, CONC group: 3 sets of 4 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: NR	18 weeks (3 sessions per week)	Increase in leg lean mass (g) with ECC (4 %), no change with CONC (2 %) or CONC (1 %), no difference between interventions

Table 2 continued

Study	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Higbie et al. 1996 [39]	ECC $n = 19$, CONC $n = 16$, CONT $n = 19$ (54F); mean age: 20.5 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 3 sets of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 60°/s	10 weeks (3 sessions per week)	Increase in quadriceps CSA (cm^2) with ECC (7 %) and CONC (5 %), no change with CONT (-1 %), greater increase with ECC vs. CONC
Hortobagyi et al. [22]	ECC $n = 7$, CONC $n = 8$, CONT $n = 6$ (21 M); mean age: 21.3 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 5.3 sets of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 60°/s	12 weeks (3 sessions per week)	Increase in quadriceps type II fibre CSA (μm^2) with ECC (38 %), no change with CONC (3 %) or CONT (NR), greater increase with ECC vs. CONC Increase in quadriceps type IIa fibre composition (%) with ECC (31 %) and CONC (22 %), no change with CONT (NR), no difference between interventions
Hortobagyi et al. [23]	ECC $n = 12$, ECC + CONC $n = 12$, CONC $n = 12$, CONT $n = 24$ (24F, 24 M); mean age: 22.0 years; training status: untrained	KE (isokinetic; single-joint)	Volume: 5.3 sets of 10 repetitions; intensity: ECC group: ECC MVC, ECC + CONC group: ECC and CONC MVC, CONC group: CONC MVC; tempo: 60°/s	12 weeks (3 sessions per week)	Increase in quadriceps type I fibre CSA (μm^2) with ECC (10 %), ECC + CONC (11 %) and CONC (4 %), no change with CONT (NR), no difference between interventions Increase in quadriceps type IIa fibre CSA (μm^2) with ECC (16 %), ECC + CONC (9 %) and CONC (5 %), no change with CONT (NR), greater increase with ECC and ECC + CONC vs. CONC Increase in quadriceps type IIx fibre CSA (μm^2) with ECC (16 %), ECC + CONC (10 %) and CONC (5 %), no change with CONT (NR), greater increase with ECC vs. ECC + CONC and CONC Increase in quadriceps muscle fibre CSA (μm^2) with ECC (52 %), no change with CONC (11 %), no difference between interventions
LaStayo et al. [36]	ECC $n = 7$, CONC $n = 7$ (14 M); mean age: 23.9 years; training status: untrained	HE and KE (isokinetic cycling; multi-joint)	Volume: 27.5 min; intensity: 62 % maximum heart rate; tempo: 60 rpm	8 weeks (3.5 sessions per week)	Increase in quadriceps type Ix fibre CSA (μm^2) with ECC (16 %), ECC + CONC (10 %) and CONC (5 %), no change with CONT (NR), greater increase with ECC vs. ECC + CONC and CONC
Malliaras et al. [26]	ECC _{Heavy} $n = 10$, ECC _{Light} $n = 10$, CONC $n = 9$, CONT $n = 9$ (38 M); mean age: 27.5 years; training status: untrained	KE (isoinertial; single-joint)	Volume: ECC _{Heavy} and CONC groups: 4 sets of 7.5 repetitions, ECC _{Light} group: 4 sets of 13.5 repetitions; intensity: ECC _{Heavy} group: 80 % ECC IRM, ECC _{Light} and CONC groups: 80 % CONC IRM; tempo: ECC: 5 s, CONC: 1 s	12 weeks (3 sessions per week)	Increase in patellar tendon Young's modulus (MPa; 50–75 %) with ECC _{Heavy} (87 %), ECC _{Light} (59 %) and CONC (81 %) vs. CONT (-3 %), no difference between interventions Increase in patellar tendon Young's modulus (MPa; 75–100 %) with ECC _{Heavy} (84 %) vs. CONC (3 %), no change with ECC _{Light} (59 %) or CONC (71 %), no difference between interventions Increase in tendon force (N) with ECC _{Heavy} (31 %) and ECC _{Light} (16 %), no change with CONC (18 %) or CONT (4 %), ECC _{Heavy} greater than CONT, no difference between interventions Increase in tendon stress (%) with ECC _{Heavy} (24 %), no change with ECC _{Light} (13 %), CONC (14 %) or CONT (2 %), no difference between interventions

Table 2 continued

Study	Population	Muscle groups (modality)	Intervention	Training duration	Training effect ($p < 0.05$)
Moore et al. [29]	ECC and CONC $n = 9$, within-subject design (9 M); mean age: 22.0 years; training status: untrained	EF (isokinetic; single-joint)	Volume: ECC group 4.44 sets of 10 repetitions, CONC group: 4.44 sets of 14 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 45°/s	9 weeks (2 sessions per week)	Increase in biceps CSA (mm^2) with ECC (7 %) and CONC (5 %), no difference between interventions
Nickols-Richardson et al. [30]	ECC $n = 33$, CONC $n = 37$ (70F); mean age: 35.0 years; training status: untrained	KF, KE, EF and EE (isokinetic; single-joint)	Volume: 4.5 sets of 6 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 60°/s	20 weeks (3 sessions per week)	Increase in lean body mass (kg) with ECC (2 %) and CONC (2 %), no difference between interventions
Rahbek et al. [73]	ECC _{whey} and ECC $n = 12$, CONC _{whey} and CONC $n = 12$, within-subject design (24 M); mean age: 23.9 years; training status: untrained	KE (isoinertial; single-joint)	Volume: 9.3 sets of 10.7 repetitions; intensity: ECC groups: 90/75 % IRM, CONC groups: 75 % IRM; tempo: ECC: 2 s, CONC: 2 s	12 weeks (2.75 sessions per week)	Increase in quadriceps CSA (cm^2) with ECC _{whey} (8 %), ECC (3 %), CONC _{whey} (6 %) and CONC (4 %), greater increase with ECC _{whey} and CONC _{whey} vs. ECC and CONC
Seger et al. [40]	ECC $n = 5$, CONC $n = 5$ (10 M); mean age: 24.5 years; training status: moderately trained	KE (isokinetic; single-joint)	Volume: 4 sets of 10 repetitions; intensity: ECC group: ECC MVC, CONC group: CONC MVC; tempo: 90°/s	10 weeks (3 sessions per week)	Increase in quadriceps CSA (cm^2) with ECC (4 %), no change with CONC (3 %), no difference between interventions
Vikne et al. [31]	ECC $n = 9$, CONC $n = 8$ (17 M); mean age: 27.1 years; training status: resistance trained	EF (isoinertial; single-joint)	Volume: 3.9 sets of 6 repetitions; intensity: ECC group: 94 % ECC IRM, CONC group: 94 % CONC IRM; tempo: ECC: 3.5 s, CONC: explosive	12 weeks (2.5 sessions per week)	Increase in biceps CSA (cm^2) with ECC (11 %), no change with CONC (3 %), greater increase with ECC vs. CONC Decrease in biceps type I fibre CSA (μm^2) with ECC (-9 %), no change with CONC (-1 %), no difference between interventions Increase in biceps type II fibre CSA (μm^2) with ECC (9 %), no change with CONC (1 %), no difference between interventions Decrease in biceps type Ix fibre composition (%) with CONC (-3 %), no change with ECC (1 %), no difference between interventions

°/s degrees per second, μm micrometre, cm centimetres, CONC concentric, CONC control, CSA cross-sectional area, ECC eccentric, ECC + CONC coupled maximal eccentric and concentric contractions, F females, kg kilograms, M males, mm millimetres, MPa megapascal, ms milliseconds, MVC maximum voluntary contraction, n number of subjects, N newtons, $N \cdot s^{-1}$ newtons per second, NR not reported, RM repetition maximum, rpm revolutions per minute, s seconds, TRAD traditional resistance training, W watts

3.3.1 Muscle Cross-Sectional Area (CSA) and Architecture

Increased muscle CSA is generally expected following a resistance training intervention of sufficient duration, and is directly related to an increase in workload and tension development [17]. The number of muscle fibres does not appear to increase during postnatal growth or as a result of training in humans [74]; however, the CSA of existing fibres increases considerably (i.e. via increased myofibril content) in response to mechanical loading [74]. Muscle hypertrophy with heavy resistance training is a product of increased protein translation, upregulation of genes involved in anabolic mechanisms, and satellite cell activation/proliferation [19, 74]. The conversion of a mechanical signal (i.e. generated during contraction) to a molecular event involves the upregulation of primary and secondary messengers within a signalling cascade to activate and/or repress pathways that regulate gene expression and protein synthesis/degradation [75]. Proteins are constantly being synthesized and broken down, even in adult muscle, therefore protein accumulation results from an increased rate of proteins being synthesized and a decreased rate of protein degradation [74]. Three factors are believed to mediate the hypertrophic signalling response to training; mechanical tension, muscle damage (i.e. exercise induced muscle damage [EIMD]) and metabolic stress [76]. The apparent superiority of eccentric training [8], or at least the inclusion of eccentric contractions (i.e. traditional training), may therefore be due to higher levels of both mechanical tension and EIMD than concentric training [77]. High levels of tension induce a mechanochemical signal to upregulate anabolic molecular and cellular activity within myofibres and satellite cells; the combined effects of active tension from contractile elements and passive tension (i.e. stretch-induced strain) from collagen content within the extracellular matrix and titin are believed to induce a more potent signal for protein synthesis [78]. Indeed, stretch-induced strain from eccentric contractions, sensed within the Z-line region of titin [79], appears to elicit a specific anabolic signalling response [75, 80]. Chronic stretch of muscle per se upregulates protein synthesis and increases the number of sarcomeres in series [74]. These observations may explain not only the increases in CSA but also the increases in fascicle length with eccentric training [11, 37, 81–85].

Franchi et al. [37] postulated that increased distal muscle hypertrophy with eccentric training reflects an increased CSA via sarcomeres in series in contrast to the addition of sarcomeres in parallel with concentric training. This mechanism would also explain the increased fascicle angle with concentric training [37], although increases in fascicle angle have also been reported following eccentric

training [11, 71, 84, 85], and it is likely that these adaptations can occur in concert to varying degrees. It is not entirely clear which mechanisms instigate the hypertrophic response in the presence of EIMD, and increases in muscle size can still occur in the absence of muscle damage [86]. Nonetheless, it is posited as an additive factor with eccentric training. The acute inflammatory response associated with eccentric contractions and EIMD is believed to induce a release of growth factors that regulate satellite cell activation/proliferation and anabolic signalling [76]. Increased cytokine activity (i.e. monocyte chemotactic protein-1 [MCP-1] and interferon gamma-induced protein 10 [IP-10]) has been reported following eccentric, but not concentric, exercise [87]. IP-10 within skeletal muscle is postulated to recruit T lymphocytes [87, 88], and T-cell infiltration has been found to enhance satellite cell activation and muscle regeneration in mice [89]. Both heavier eccentric loading and fast eccentric contractions involve high levels of tension generation per active motor unit [90, 91], and influence the magnitude of EIMD [92, 93], which may explain the enhanced hypertrophic response when these two variables are emphasized [16, 17, 94]. A greater resistance to fatigue has been observed across a series of maximal contractions [95], and it is probable that metabolic stress will be lower during eccentric versus concentric training [96]. Interestingly, the utilization of extended-duration eccentric cycling (i.e. 20–30 min) of a relatively low contraction intensity per repetition has been demonstrated to lead to increases in muscle CSA [34, 36]. It is possible that under these conditions there is a progressive recruitment and fatigue of higher threshold motor units, which in turn can instigate a marked hypertrophic response independent of high levels of mechanical tension [97]. Therefore, the inclusion of concentric contractions (i.e. traditional resistance training), eccentric contractions to fatigue, or a combination of eccentric loading and blood-flow restriction [98], may be warranted if maximal hypertrophy is the objective.

3.3.2 Muscle Fibre Size and Composition

Greater hypertrophy in type II muscle fibres compared with type I muscle fibres is not uncommon with resistance training [99]. Fast contracting fibres are recruited infrequently and their selective hypertrophy is proposed to be an adaptive response to aid power development during periods of near-maximal recruitment [74]. A greater increase in type II fibre area when comparing eccentric with concentric training may reflect the overall greater increase in muscle CSA. It has been further suggested that increased recruitment, tension-generating capacity and predisposition to damage of fast-twitch fibres [81] contribute to the propensity for type IIa [22, 94] and IIx [19, 94] fibre

hypertrophy with eccentric training. Eccentric training appears to exert a unique influence on the MHC phenotype shift with either improved maintenance of [13, 31, 100] or increase in [101] IIX fibre composition following training. A shift in fibre type composition (i.e. MHC isoform), particularly a shift from MHC IIX to IIA is commonly reported following resistance training [22, 31, 102], which can be subsequently reversed, or overshoot above pre-existing levels, with detraining [103]. Increased muscle activity via either endurance or traditional resistance exercise appears to switch off the IIX gene in IIX fibres, thus increasing the proportion of IIA fibres at the expense of IIX fibres [102]. It is proposed that both the total number of contractions (i.e. nerve impulses) and maximal tensile load exerted on muscle mediate the MHC shift [102]. The MHC IIX reduction and subsequent overshoot phenomena are not well understood, but it has been postulated that IIX is the 'default' MHC gene [102, 104], and training appears to induce a shift towards a more fatigue-resistant phenotype [103]. Multiple lines of evidence are suggestive of a specific response with eccentric contractions. Friedmann-Bette and colleagues found an increase in IIA in situ hybridizations, or fibres expressing elevated levels of IIX messenger RNA (mRNA), which were postulated to be in transition [19], and a tendency towards an increase in IIX mRNA has also been reported following a submaximal eccentric training protocol [105]. While changes in fibre type and MHC mRNA do not always occur in parallel, long-term steady state levels of mRNA are reasonably well correlated with muscle fibre composition [105]. Furthermore, a 10-week, fast (i.e. 180°/s) isokinetic eccentric training intervention of the elbow flexors was demonstrated to increase type IIX composition by 7 % [101]. These chronic findings align with observations of acute satellite cell [106] and anabolic signalling pathway upregulation [107] in type II fibres following eccentric exercise, and support the proposition of a shift towards a faster phenotype with eccentric training [6].

3.3.3 Tendon CSA and Qualitative Properties

Historically, tendon tissue was considered inert, relatively non-vascular and inelastic. However, recent research has highlighted the dynamic nature of the extracellular matrix, and thus adaptive capacity in response to mechanical load [108]. Changes in tendon structure and function appear to be partly mediated by contraction type [26]. Unfortunately, few studies have compared the effects of eccentric training with other resistance training modalities on tendon adaptation. Eccentric loading has gained widespread implementation within the rehabilitation setting as a tool to manage lower limb tendinopathy and improve tendon structure [108, 109]. However, the mechanisms

underpinning the improvements with eccentric loading are not well understood [26]. Chronic loading has been demonstrated to increase tendon stiffness, and possibly CSA, and it seems maximizing tendon strain may be necessary in optimizing the adaptive response [26, 85, 110]. In one investigation comparing heavy eccentric, light eccentric and concentric training, only heavy eccentric loading elicited improvements in patellar tendon Young's modulus (75–100 % MVC) and strain (%) without concomitant increases in tendon CSA [26], aligning with a previous report of heavy eccentric training inducing a reduction in elbow flexor series elastic component compliance [111]. In contrast to these findings, a decrease [112], or no change [108], in Achilles tendon stiffness has been reported following the implementation of a popular submaximal eccentric heel drop protocol [113]. These observations may be related to the contraction intensity, and thus provide further support for the importance of load for increasing tendon stiffness. Changes in tendon stiffness in the absence of increases in CSA may be due to increased collagen packing density, alterations in crimp angle, or increased water content [26, 114]. As the tendon is metabolically active [112, 114], adaptation may be driven by changes in rates of protein synthesis and degradation similar to muscle tissue in a coordinated musculo-tendinous response [115]. The cellular tensegrity model proposes that cells (i.e. myofibres and fibroblasts) can respond, in a coordinated fashion, to mechanical stress via integrins located in the plasma membrane and proteins connecting the extracellular matrix to the cytoskeleton, which stimulates functional remodelling of the MTU [115]. The magnitude of mechanical stress may augment the signalling response and thus explain the greater adaptation with heavy versus light eccentric training [26]. The finding that eccentric training can increase tendon CSA [72] is of interest as other acute interventions seem to be less effective at eliciting quantitative changes in tendon tissue. Plyometric training [116] and traditional resistance training [117] have been found to increase Achilles tendon stiffness, but not CSA. Long-term endurance training is associated with increased Achilles tendon CSA [118], which suggests that long-term chronic loading may be necessary to elicit quantitative changes in tendon with submaximal loads. It appears that heavy eccentric training can induce both qualitative and quantitative changes in tendon, although more research is necessary to clarify the optimal loading conditions.

3.3.4 Summary of Eccentric Training Effects on Muscle-Tendon Unit Morphology and Architecture

Eccentric training appears to elicit greater increases in muscle CSA than concentric or traditional resistance training. The combination of heavier absolute loads and a

smaller number of recruited motor units during eccentric training [52] involves high levels of mechanical tension per motor unit [91] and a greater propensity for EIMD [119]. These factors, combined with stretch-induced strain [78], may stimulate the hypertrophic signalling response to a greater extent than concentric or traditional resistance training [76]. Heavy eccentric loads and fast contraction velocities both appear to further stimulate the adaptive response [16, 17, 94, 101]. The pattern of increased muscle CSA appears to be mediated by contraction type, and eccentric training may promote the addition of sarcomeres in series, as inferred from changes in muscle fascicle length [11, 37]. Selective increases in fast-twitch fibre size have been reported and there is evidence to suggest that a shift towards a fast phenotype can occur as a result of chronic eccentric training [6]. The predisposition for fast-twitch fibre hypertrophy probably results from an increased recruitment with heavy loads, the tension generating capacity and subsequent damage of these fibres [19, 22]. While further research is required, it is possible an upregulation of IIX mRNA signalling can occur with eccentric training [19, 105], leading to an attenuated IIX to IIA shift [13, 31, 100], or an increase in IIX composition with fast contractions [101]. Eccentric training may promote increases in tendon stiffness [26] and CSA [72], which influence the storage and return of elastic strain energy and probably contribute to the observed enhancements in SSC performance. The MTU morphological and architectural adaptations, in conjunction with changes in muscle mechanical function (Sect. 3.2), have important implications for strength, power and speed performance; however, more research is necessary to determine how these findings translate to trained athletes.

4 Conclusions

Eccentric training using external loads greater than the relative concentric training intensity is a potent stimulus for enhancements in muscle mechanical function, and MTU morphological and architectural adaptations. The inclusion of eccentric loads above maximal concentric strength is therefore an avenue to induce novel training stimuli and effect change in key determinants, and functional metrics, of strength, power and speed performance. Strength improvements are largely mode-specific and arise from a combination of neural, morphological and architectural adaptations [8]. Increased agonist volitional drive is posited as the primary contributing factor to the marked increases in eccentric strength observed following training [54]. Eccentric training improves concentric muscle power and SSC performance to a greater extent than concentric or traditional modalities [13, 34, 46, 47]. Reports on changes

in RFD have been mixed [27, 33], although limited research in this area has been undertaken. Improvements in muscle power and SSC performance are likely related to improvements in total strength, an improved ability to rapidly recruit fast motor units, qualitative changes to tendon tissue, and an enhanced eccentric phase within the SSC [3]. Eccentric training can elicit greater increases in muscle CSA than concentric or traditional resistance training. High levels of mechanical tension per active motor unit [91], stretch-induced strain [78], and a greater propensity for EIMD [119] with eccentric training may stimulate anabolic signalling to a greater extent than concentric or traditional resistance training [76]. The nature of hypertrophy appears to differ with eccentric training versus concentric training, and the addition of sarcomeres in series, as inferred from changes in muscle fascicle length, may contribute to increases in CSA [11, 37]. A greater number of sarcomeres in series may subsequently increase muscle shortening velocity and increase force production at longer muscle lengths [6]. Fast-twitch fibres hypertrophy to a greater extent as a consequence of eccentric training [19, 22], and while further research is required, it is possible an upregulation of IIX mRNA signalling can occur [19, 105], leading to an attenuated IIX to IIA shift [13, 31, 100], or an increase in IIX composition with fast contractions [101]. An increased number of sarcomeres in series, selective fast-twitch fibre hypertrophy, and possibly increased IIX composition, have been collectively referred to as a shift towards a faster phenotype [6] and may contribute to improvements in speed and power performance with eccentric training. Furthermore, increases in tendon stiffness [26] and CSA [72] will aid the storage and return of elastic strain energy during SSC movements. Particular consideration should be given to contraction velocity; fast eccentric training appears to induce the largest improvements in strength, power and SSC performance [17, 47], and further stimulates increases in muscle CSA [17] and IIX fibre composition [101]. While fast contractions per se are probably an important stimulus, further research is necessary in elucidating the influence of a markedly lower time under tension with such protocols [94]. An acute eccentric training intervention in untrained participants seems to follow the typical adaptive pattern with resistance training. The largest increases in early-phase strength (i.e. 3–4 weeks) are underpinned by neural factors, while MTU morphological and architectural changes require a longer time to materialize [84]. As residual fatigue with eccentric exercise can suppress force production and affect neural control for a period of time following the cessation of training, an appropriate recovery window (i.e. up to 8 weeks) may be necessary to fully realize neuromuscular adaptations [71]. It is less clear how these findings translate to resistance-trained subjects as the majority of

investigations have recruited untrained participants; it has been suggested that the pattern of adaptation may be similar, but of a lower magnitude [6]. The heterogeneity of protocols used makes it difficult to elucidate best practice in training volume and frequency. Presently, it would seem that the management of EIMD and delayed-onset muscle soreness should be a primary consideration. Further research is necessary to determine the optimal progression of load and contraction velocity, especially in resistance-trained subjects, for the MTU and performance outcomes.

Compliance with Ethical Standards

Funding No sources of funding were used to assist in the preparation of this article.

Conflict of interest Jamie Douglas, Simon Pearson, Angus Ross and Mike McGuigan declare that they have no conflicts of interest relevant to the content of this review.

References

- McGuigan MR, Wright GA, Fleck SJ. Strength training for athletes: does it really help sports performance? *Int J Sports Physiol Perform.* 2012;7:2–5.
- Beattie K, Kenny IC, Lyons M, et al. The effects of strength training on performance in endurance athletes. *Sports Med.* 2014;44:845–65.
- Cormie P, McGuigan MR, Newton RU. Developing maximal neuromuscular power: part 1. Biological basis of maximal power production. *Sports Med.* 2011;41(1):17–38.
- Lindstedt SL, LaStayo PC, Reich TE. When active muscles lengthen: properties and consequences of eccentric contractions. *News Physiol Sci.* 2001;16:256–61.
- Hortobágyi T, Katch F. Eccentric and concentric torque velocity relationships during arm flexion and extension: influence of strength level. *Eur J Appl Physiol.* 1990;60:395–401.
- Vogt M, Hoppeler HH. Eccentric exercise: mechanisms and effects when used as training regime or training adjunct. *J Appl Physiol (1985).* 2014;116(11):1446–54.
- Isner-Horobeti M, Dufour SP, Vautravers P, et al. Eccentric exercise training: modalities, applications and perspectives. *Sports Med.* 2013;43:483–512.
- Roig M, O'Brien K, Kirk G, et al. The effects of eccentric versus concentric resistance training on muscle strength and mass in healthy adults: a systematic review with meta-analysis. *Br J Sports Med.* 2009;43(8):556–68.
- Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med.* 2009;151(4):264–9.
- Ben-Sira D, Ayalon A, Tavi M. The effect of different types of strength training on concentric strength in women. *J Strength Cond Res.* 1995;9(3):143–8.
- Blazevich AJ, Cannavan D, Coleman DR, et al. Influence of concentric and eccentric resistance training on architectural adaptation in human quadriceps muscles. *J Appl Physiol (1985).* 2007;103(5):1565–75.
- Brandenburg JP, Docherty D. The effects of accentuated eccentric loading on strength, muscle hypertrophy, and neural adaptations in trained individuals. *J Strength Cond Res.* 2002;16(1):25–32.
- Colliander EB, Tesch PA. Effects of eccentric and concentric muscle actions in resistance training. *Acta Physiol Scand.* 1990;140(1):31–9.
- Colliander EB, Tesch PA. Effects of detraining following short term resistance training on eccentric and concentric muscle strength. *Acta Physiol Scand.* 1992;144(1):23–9.
- Ellenbecker TS, Davies GJ, Rowinski MJ. Concentric versus eccentric isokinetic strengthening of the rotator cuff. Objective data versus functional test. *Am J Sports Med.* 1988;16(1):64–9.
- English K, Loehr J, Lee S, et al. Early-phase musculoskeletal adaptations to different levels of eccentric resistance after 8 weeks of lower body training. *Eur J Appl Physiol.* 2014;114(11):2263–80.
- Farthing JP, Chilibeck PD. The effects of eccentric and concentric training at different velocities on muscle hypertrophy. *Eur J Appl Physiol.* 2003;89:578–86.
- Farup J, Rahbek SK, Riis S, et al. Influence of exercise contraction mode and protein supplementation on human skeletal muscle satellite cell content and muscle fiber growth. *J Appl Physiol.* 1985;2014(117):898–909.
- Friedmann-Bette B, Bauer T, Kinscherf R, et al. Effects of strength training with eccentric overload on muscle adaptation in male athletes. *Eur J Appl Physiol.* 2010;108(4):821–36.
- Godard MP, Wygand JW, Carpinelli RN, et al. Effects of accentuated eccentric resistance training on concentric knee extensor strength. *J Strength Cond Res.* 1998;12(1):26–9.
- Hawkins SA, Schroeder ET, Wiswell RA, et al. Eccentric muscle action increases site-specific osteogenic response. *Med Sci Sports Exerc.* 1999;31(9):1287–92.
- Hortobágyi T, Hill JP, Houmar JA, et al. Adaptive responses to muscle lengthening and shortening in humans. *J Appl Physiol (1985).* 1996;80(3):765–72.
- Hortobágyi T, Dempsey L, Fraser D, et al. Changes in muscle strength, muscle fibre size and myofibrillar gene expression after immobilization and retraining in humans. *J Physiol.* 2000;524(Pt 1):293–304.
- Kaminski TW, Wabbersen CV, Murphy RM. Concentric versus enhanced eccentric hamstring strength training: clinical implications. *J Athletic Train.* 1998;33(3):216–21.
- Komi PV, Buskirk ER. Effect of eccentric and concentric muscle conditioning on tension and electrical activity of human muscle. *Ergonomics.* 1972;15(4):417–34.
- Malliaras P, Kamal B, Nowell A, et al. Patellar tendon adaptation in relation to load-intensity and contraction type. *J Biomech.* 2013;46(11):1893–9.
- Miller LE, Pierson LM, Nickols-Richardson SM, et al. Knee extensor and flexor torque development with concentric and eccentric isokinetic training. *Res Q Exerc Sport.* 2006;77(1):58–63.
- Mont MA, Cohen DB, Campbell KR, et al. Isokinetic concentric versus eccentric training of shoulder rotators with functional evaluation of performance enhancement in elite tennis players. *Am J Sports Med.* 1994;22(4):513–7.
- Moore D, Young M, Phillips S. Similar increases in muscle size and strength in young men after training with maximal shortening or lengthening contractions when matched for total work. *Eur J Appl Physiol.* 2012;112(4):1587–92.
- Nickols-Richardson SM, Miller LE, Wootten DF, et al. Concentric and eccentric isokinetic resistance training similarly increases muscular strength, fat-free soft tissue mass, and specific bone mineral measurements in young women. *Osteoporos Int.* 2007;18(6):789–96.
- Vikne H, Refsnes PE, Ekmark M, et al. Muscular performance after concentric and eccentric exercise in trained men. *Med Sci Sports Exerc.* 2006;38(10):1770–81.

32. Yarrow JF, Borsa PA, Borst SE, et al. Early-phase neuroendocrine responses and strength adaptations following eccentric-enhanced resistance training. *J Strength Cond Res.* 2008;22(4):1205–14.
33. Blazevich AJ, Horne S, Cannavan D, et al. Effect of contraction mode of slow-speed resistance training on the maximum rate of force development in the human quadriceps. *Muscle Nerve.* 2008;38:1133–46.
34. Gross M, Lüthy F, Kroell J, et al. Effects of eccentric cycle ergometry in alpine skiers. *Int J Sports Med.* 2010;31(8):572–6.
35. LaStayo PC, Reich TE, Urquhart M, et al. Chronic eccentric exercise: improvements in muscle strength can occur with little demand for oxygen. *Am J Physiol.* 1999;276(2):R611–5.
36. LaStayo PC, Pierotti DJ, Pifer J, et al. Eccentric ergometry: increases in locomotor muscle size and strength at low training intensities. *Am J Physiol.* 2000;278(5):R1282–8.
37. Franchi MV, Atherton PJ, Reeves ND, et al. Architectural, functional and molecular responses to concentric and eccentric loading in human skeletal muscle. *Acta Physiol.* 2014;210:642–54.
38. Duncan PW, Chandler JM, Cavanaugh DK, et al. Mode and speed specificity of eccentric and concentric exercise training. *J Orthop Sports Phys Ther.* 1989;11(2):70–5.
39. Higbie EJ, Cureton KJ, Warren GL, et al. Effects of concentric and eccentric training on muscle strength, cross-sectional area, and neural activation. *J Appl Physiol* (1985). 1996;81(5):2173–81.
40. Seger JY, Arvidsson B, Thorstensson A. Specific effects of eccentric and concentric training on muscle strength and morphology in humans. *Eur J Appl Physiol.* 1998;79(1):49–57.
41. Tomberlin JP, Basford JR, Schwen EE, et al. Comparative study of isokinetic eccentric and concentric quadriceps training. *J Orthop Sports Phys Ther.* 1991;14(1):31–6.
42. Spurway NC. The effect of strength training on the apparent inhibition of eccentric force production in voluntary activated human quadriceps. *Eur J Appl Physiol.* 2000;82:374–80.
43. Barstow IK, Bishop MD, Kaminski TW. Is enhanced-eccentric resistance training superior to traditional training for increasing elbow flexor strength? *J Sport Sci Med.* 2003;2:62–9.
44. Farthing JP, Chilibeck PD. The effect of eccentric training at different velocities on cross-education. *Eur J Appl Physiol.* 2003;89(6):570–7.
45. Hortobagyi T, Lambert NJ, Hill JP. Greater cross education following training with muscle lengthening than shortening. *Med Sci Sports Exerc.* 1997;29(1):107–12.
46. Elmer S, Hahn S, McAllister P, et al. Improvements in multi-joint leg function following chronic eccentric exercise. *Scand J Med Sci Sports.* 2012;22(5):653–61.
47. Liu C, Chen CS, Ho WH, et al. The effects of passive leg press training on jumping performance, speed, and muscle power. *J Strength Cond Res.* 2013;27(6):1479–86.
48. Aagaard P, Simonsen EB, Andersen JL, et al. Neural inhibition during maximal eccentric and concentric quadriceps contraction: effects of resistance training. *J Appl Physiol.* 1985;2000(89):2249–57.
49. Vangsgaard S, Taylor JL, Hansen EA, et al. Changes in H reflex and neuromechanical properties of the trapezius muscle after 5 weeks of eccentric training: a randomized controlled trial. *J Appl Physiol.* 1985;2014(116):1623–31.
50. Pensini M, Martin A, Maffiuletti NA. Central versus peripheral adaptations following eccentric resistance training. *Int J Sports Med.* 2002;23:567–74.
51. Beltman JGM, Sargeant AJ, van Mechelen W, et al. Voluntary activation level and muscle fiber recruitment of human quadriceps during lengthening contractions. *J Appl Physiol* (1985). 2004;97(2):619–26.
52. Duchateau J, Baudry S. Insights into the neural control of eccentric contractions. *J Appl Physiol* (1985). 2014;116(11):1418–25.
53. Duclay J, Martin A, Robbe A, et al. Spinal reflex plasticity during maximal dynamic contractions after eccentric training. *Med Sci Sports Exerc.* 2008;40(4):722–34.
54. Aagaard P. Training-induced changes in neural function. *Exerc Sport Sci Rev.* 2003;31(2):61–7.
55. Sale DG. Neural adaptation to resistance training. *Med Sci Sports Exerc.* 1988;20(5):S135–45.
56. Aagaard P. Neural adaptations to resistance exercise. In: Cardinale M, Newton R, Nosaka K, editors. *Strength and conditioning: biological principles and practical applications.* Chichester: Wiley-Blackwell; 2011. p. 105–24.
57. Van Cutsem M, Duchateau J, Hainaut K. Changes in single motor unit behaviour contribute to the increase in contraction speed after dynamic training in humans. *J Physiol.* 1998;513(1):295–305.
58. Duchateau J, Semmler JG, Enoka RM. Training adaptations in the behaviour of human motor units. *J Appl Physiol.* 1985;2006(101):1766–75.
59. Papadopoulos C, Theodosiou K, Bogdanis GC, et al. Multiarticular isokinetic high-load eccentric training induces large increases in eccentric and concentric strength and jumping performance. *J Strength Cond Res.* 2014;28(9):2680–8.
60. Cormie P, McGuigan MR, Newton RU. Changes in the eccentric phase contribute to improved stretch-shorten cycle performance after training. *Med Sci Sports Exerc.* 2010;42(9):1731–44.
61. Cook CJ, Beaven CM, Kilduff LP. Three weeks of eccentric training combined with overspeed exercises enhances power and running speed performance gains in trained athletes. *J Strength Cond Res.* 2013;27(5):1280–6.
62. de Hoyo M, Pozzo M, Sanudo B, et al. Effects of a 10-week in-season eccentric-overload training program on muscle-injury prevention and performance in junior elite soccer players. *Int J Sports Physiol Perform.* 2015;10:46–52.
63. Bojsen-Moller J, Magnusson SP, Rasmussen LR, et al. Muscle performance during maximal isometric and dynamic contractions is influenced by the stiffness of the tendinous structures. *J Appl Physiol.* 1985;2005(99):986–94.
64. Andersen LL, Aagaard P. Influence of maximal muscle strength and intrinsic muscle contractile properties on contractile rate of force development. *Eur J Appl Physiol.* 2006;96:46–52.
65. Oliveira AS, Corvino RB, Caputo F, et al. Effects of fast-velocity eccentric resistance training on early and late rate of force development. *Eur J Sport Sci.* 2016;16(2):199–205.
66. Bottinelli R, Canepari M, Pellegrino MA, et al. Force-velocity properties of human skeletal muscle fibres: myosin heavy chain isoform and temperature dependence. *J Physiol.* 1996;495(2):573–86.
67. Bottinelli R, Pellegrino MA, Canepari M, et al. Specific contributions of various muscle fibre types to human muscle performance: an in vitro study. *J Electromyogr Kinesiol.* 1999;9:87–95.
68. Andersen LL, Andersen JL, Zebis MK, et al. Early and late rate of force development: differential adaptive responses to resistance training? *Scand J Med Sci Sports.* 2010;20:162–9.
69. de Oliveira FBD, Rizzato GF, Denadai BS. Are early and late rate of force development differently influenced by fast-velocity resistance training. *Clin Physiol Funct Imaging.* 2013;33:282–7.
70. Tillin NA, Pain MTG, Folland JP. Short-term training for explosive strength causes neural and mechanical adaptations. *Exp Physiol.* 2012;97(5):630–41.
71. Leong CH, McDermott WJ, Elmer SJ, et al. Chronic eccentric cycling improves quadriceps muscle structure and maximum cycling power. *Int J Sports Med.* 2014;35:559–65.

72. Farup J, Rahbek SK, Vendelbo MH, et al. Whey protein hydrolysate augments tendon and muscle hypertrophy independent of resistance exercise contraction mode. *Scand J Med Sci Sports*. 2014;24:788–98.
73. Rahbek SK, Farup J, Moller AB, et al. Effects of divergent resistance exercise contraction mode and dietary supplementation type on anabolic signalling, muscle protein synthesis and muscle hypertrophy. *Amino Acids*. 2014;46(10):2377–92.
74. Goldspink G, Harridge S. Cellular and molecular aspects of adaptation in skeletal muscle. In: Komi PV, editor. *Strength and power in sport*. Volume III encyclopaedia of sports medicine. Encyclopaedia of Sports Medicine. Osney Mead: Blackwell Science Ltd; 2003. p. 231–51.
75. Coffey VG, Hawley JA. The molecular bases of training adaptation. *Sports Med*. 2007;37(9):737–63.
76. Schoenfeld BJ. The mechanisms of muscle hypertrophy and their application to resistance training. *J Strength Cond Res*. 2010;24(10):2857–72.
77. Eliasson J, Elfegoun T, Nilsson J, et al. Maximal lengthening contractions increase p70 S6 kinase phosphorylation in human skeletal muscle in the absence of nutritional supply. *Am J Physiol Endocrinol Metab*. 2006;291:E1197–205.
78. Toigo M, Boutellier U. New fundamental resistance exercise determinants of molecular and cellular muscle adaptations. *Eur J Appl Physiol*. 2006;97:643–63.
79. Miller MK, Bang ML, Witt CC, et al. The muscle ankyrin repeat proteins: CARP, ankrd2/Arpp and DARP as a family of titin filament-based stress response molecules. *J Mol Biol*. 2003;333:951–64.
80. Kumar A, Chaudhry I, Reid MB, et al. Distinct signalling pathways are activated in response to mechanical stress applied axially and transversely to skeletal muscle fibers. *J Biol Chem*. 2002;277(48):46493–503.
81. Proske U, Morgan DL. Muscle damage from eccentric exercise: mechanism, mechanical signs, adaptation and clinical adaptations. *J Physiol*. 2001;537:333–45.
82. Baroni BM, Geremia JM, Rodrigues R, et al. Muscle architecture adaptations to knee extensor eccentric training: rectus femoris vs. vastus lateralis. *Muscle Nerve*. 2013;48:498–506.
83. Potier TG, Alexander CM, Seynnes OR. Effects of eccentric strength training on biceps femoris muscle architecture and knee joint range of movement. *Eur J Appl Physiol*. 2009;105:939–44.
84. Seynnes OR, de Boer M, Narici MV. Early skeletal muscle hypertrophy and architectural changes in response to high-intensity resistance training. *J Appl Physiol*. 1985;2007(102):368–73.
85. Duclay J, Martin A, Duclay A, et al. Behavior of fascicles and the myotendinous junction of human medial gastrocnemius following eccentric strength training. *Muscle Nerve*. 2009;39:819–27.
86. Schoenfeld BJ. Does exercise-induced muscle damage play a role in skeletal muscle hypertrophy? *J Strength Cond Res*. 2012;26(5):1441–53.
87. Hyldahl RD, Olson T, Welling T, et al. Satellite cell activity is differentially affected by contraction mode in human muscle following a work-matched bout of exercise. *Front Physiol*. 2014;5(485):1–11.
88. Crescioli C, Sottili M, Bonini P, et al. Inflammatory response in human skeletal muscle cells: CXCL10 as a potential therapeutic agent. *Eur J Cell Biol*. 2012;91:139–49.
89. Burzyn D, Kuswanto W, Kolodin D, et al. A special population of regulatory T cells potentiates muscle repair. *Cell*. 2013;155(6):1282–95.
90. Hollander DB, Kraemer RR, Kilpatrick MW, et al. Maximal eccentric and concentric strength discrepancies between young men and women for dynamic resistance exercise. *J Strength Cond Res*. 2007;21(3):34–40.
91. Prilutsky BI. Eccentric muscle action in sport and exercise. In: Zatsiorsky VM, editor. *Biomechanics in sport*. Volume IX encyclopaedia of sports medicine. Encyclopaedia of Sports Medicine. Osney Mead: Blackwell Science Ltd; 2000. p. 56–86.
92. Chapman D, Newton M, Sacco P, et al. Greater muscle damage induced by fast versus slow velocity eccentric exercise. *Int J Sports Med*. 2006;27:591–8.
93. McHugh MP, Tetro DT. Changes in the relationship between joint angle and torque production associated with the repeated bout effect. *J Sport Sci*. 2003;21:927–32.
94. Shepstone TN, Tang JE, Dallaire S, et al. Short-term high- vs. low-velocity isokinetic lengthening training results in greater hypertrophy of the elbow flexors in young men. *J Appl Physiol*. 1985;2005(98):1768–76.
95. Baroni BM, Stocchero CMA, do Espírito Santo RC, et al. The effect of contraction type on muscle strength, work and fatigue in maximal isokinetic exercise. *Isokinet Exerc Sci*. 2011;19(3):215–20.
96. Horstmann T, Mayer F, Maschmann J, et al. Metabolic reaction after concentric and eccentric endurance-exercise of the knee and ankle. *Med Sci Sports Exerc*. 2001;33(5):791–5.
97. Morton RW, Oikawa SY, Wavell CG, et al. Neither load nor systemic hormones determine resistance training-mediated hypertrophy or strength gains in resistance-trained young men. *J Appl Physiol*. 1985;2016(121):129–38.
98. Sudo M, Ando S, Poole DC, et al. Blood flow restriction prevents muscle damage but not protein synthesis signalling following eccentric contractions. *Physiol Rep*. 2015;3(7):1–10.
99. Campos GER, Luecke TJ, Wendeln HK, et al. Muscular adaptations in response to three different resistance-training regimens: specificity of repetition maximum training zones. *Eur J Appl Physiol*. 2002;88:50–60.
100. Raue U, Terpstra B, Williamson DL, et al. Effects of short-term concentric vs. eccentric resistance training on single muscle fiber MHC distribution in humans. *Int J Sports Med*. 2005;26:339–43.
101. Paddon-Jones D, Leveritt M, Lonergan A, et al. Adaptation to chronic eccentric exercise in humans: the influence of contraction velocity. *Eur J Appl Physiol*. 2001;85:466–71.
102. Andersen JL, Aagaard P. Myosin heavy chain IIX overshoot in human skeletal muscle. *Muscle Nerve*. 2000;23:1095–104.
103. Staron RS, Johnson P. Myosin polymorphism and differential expression in adult human skeletal muscle. *Comp Biochem Physiol*. 1993;106B(3):463–75.
104. Andersen JL, Mohr T, Biering-Sorensen F, et al. Myosin heavy chain isoform transformation in single fibres from m. vastus lateralis in spinal cord injured individuals: effects of long-term functional electrical stimulation (FES). *Pflügers Arch*. 1996;431:513–8.
105. Friedmann B, Kinscherf R, Vorwald S, et al. Muscular adaptations to computer-guided strength training with eccentric overload. *Acta Physiol Scand*. 2004;182:77–88.
106. Cermak NM, Snijders T, McKay BR, et al. Eccentric exercise increases satellite cell content in type II muscle fibers. *Med Sci Sports Exerc*. 2013;45(2):230–7.
107. Tannerstedt J, Apró W, Blomstrand E. Maximal lengthening contractions induce different signalling responses in the type I and type II fibers of human skeletal muscle. *J Appl Physiol*. 1985;2009(106):1412–8.
108. Mahieu NN, McNair P, Cools A, et al. Effects of eccentric training on the plantar flexor muscle-tendon tissue properties. *Med Sci Sports Exerc*. 2008;40(1):117–23.
109. Ohberg L, Lorentzon R, Alfredson H. Eccentric training in patients with chronic Achilles tendinosis: normalised tendon

- structure and decreased thickness at follow up. *Br J Sports Med.* 2004;38:8–11.
110. Arampatzis A, Karamanidis K, Albracht K. Adaptational responses of the human Achilles tendon by modulation of the applied cyclic strain magnitude. *J Exp Biol.* 2007;210:2743–53.
 111. Pousson M, Van Hoecke J, Goubel F. Changes in elastic characteristics of human muscle induced by eccentric exercise. *J Biomech.* 1990;23(4):343–8.
 112. Morrissey D, Roskilly A, Twycross-Lewis R, et al. The effect of eccentric and concentric calf muscle training on Achilles tendon stiffness. *Clin Rehabil.* 2011;25:238–47.
 113. Alfredson H, Pietila T, Jonsson P, et al. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med.* 1998;26(3):360–6.
 114. Magnusson SP, Narici MV, Maganaris CN, et al. Human tendon behaviour and adaptation, in vivo. *J Physiol.* 2008;586(1):71–81.
 115. Miller BF, Olesen JL, Hansen M, et al. Coordinated collagen and muscle protein synthesis in human patella tendon and quadriceps muscle after exercise. *J Physiol.* 2005;567(3):1021–33.
 116. Foure A, Nordez A, Cornu C. Plyometric training effects on Achilles tendon stiffness and dissipative properties. *J Appl Physiol.* 1985;2010(109):849–54.
 117. Kubo K, Morimoto M, Komuro T, et al. Effects of plyometric and weight training on muscle-tendon complex and jump performance. *Med Sci Sports Exerc.* 2007;39(10):1801–10.
 118. Magnusson SP, Kjaer M. Region-specific differences in Achilles tendon cross-sectional area in runners and non-runners. *Eur J Appl Physiol.* 2003;90:549–53.
 119. McHugh MP. Recent advances in the understanding of the repeated bout effect: the protective effect against muscle damage from a single bout of eccentric exercise. *Scand J Med Sci Sports.* 2003;13(2):88–97.